

Einstein Gravity in Almost Kähler and Lagrange–Finsler Variables and Deformation Quantization

Sergiu I. Vacaru*

*Science Department, University "Al. I. Cuza" Iași,
54 Lascar Catargi street, 700107, Iași, Romania*

April 5, 2010

Abstract

A geometric procedure is elaborated for transforming (pseudo) Riemannian metrics and connections into canonical geometric objects (metric and nonlinear and linear connections) for effective Lagrange, or Finsler, geometries which, in turn, can be equivalently represented as almost Kähler spaces. This allows us to formulate an approach to quantum gravity following standard methods of deformation quantization. Such constructions are performed not on tangent bundles, as in usual Finsler geometry, but on spacetimes enabled with nonholonomic distributions defining 2+2 splitting with associate nonlinear connection structure. We also show how the Einstein equations can be written in terms of Lagrange–Finsler variables and corresponding almost symplectic structures and encoded into the zero-degree cohomology coefficient for a quantum model of Einstein manifolds.

Keywords: Einstein spaces, Lagrange geometry, Finsler geometry, deformation quantization, quantum gravity

MSC: 83C99, 53D55, 53B40, 53B35

PACS: 04.20.-q, 02.40.-k, 02.90.+g, 02.40.Yy

*e-mails: sergiu.vacaru@uaic.ro, Sergiu.Vacaru@gmail.com,
<http://www.scribd.com/people/view/1455460-sergiu>

1 Introduction

The ideas presented in this article grew out of our attempts [1, 2, 3, 4] to reformulate equivalently the Einstein gravity and generalizations as almost Kähler geometries in order to apply the deformation quantization methods [5, 6, 7, 8] and elaborate quantum models of such spaces. The term "quantization" usually means a procedure stating a quantum model for a given classical theory. One follows certain analogy with existing approaches to quantum and classical mechanics and field theories. Such constructions involve a great amount of ambiguity because quantum theories provide more refined descriptions of physical systems than the classical ones. For gravity and gauge theories, this is related to a set of unsolved yet fundamental mathematical problems in nonlinear functional analysis and noncommutative geometry.

A deeper level of description of quantum systems connected to a consistent quantization formalism manifests itself in extra geometric structures to be defined on a generalized phase space. For instance, a symplectic connection and related Poisson structures, which are not restricted to classical dynamics, become key ingredients of geometric quantum theories. The existing examples of deformation quantization of gravity and matter fields involve more "rigid" geometric structures (like metric and nonlinear and linear connections, torsions, almost complex structures etc). Different choices for such structures lead, in general, to inequivalent quantizations on the same phase space, a typical feature of nonlinear theories.

This work was stimulated in part by many attempts to apply deformation quantization to generalized Poisson structures, gravity and string theory. Here we mention some approaches and emphasize the following key ideas and results:

A formal scheme proposing a unification of the four dimensional Einstein gravity and quantum mechanics was proposed, which resulted in loop quantum gravity (LQG) and spin network theory, see comprehensive reviews in [9, 10, 11]. We also cite a discussion [12] on alternative approaches following methods which differ from those proposed in string theory [13, 14, 15], which aim to formulate a unified higher dimensional theory of interactions. The background free LQG originated from a canonical formulation of general relativity, based on ADM 3+1 decomposition and the Palatini formalism (see a review of results in [16]), and Ashtekar's connection dynamics with a relatively simple Hamiltonian constraint and further modifications.

However, the attempts to quantize nonlinear field theories, including different models of gravity, were oriented not only to the canonical quantum

gravity and/or LQG but also to alternative methods of geometric quantization. The fundamental works [17, 18, 19] preceding the Fedosov and Kontsevich approaches [5, 6, 7, 8] should also be mentioned. In Ref. [20], a deformation quantization for self-dual gravity formulated in Plebański (self-dual) variables was considered. A variant of deformation (Moyal) quantization for general constrained Hamiltonian systems was introduced in [21]. It was shown in [21] how second class constraints can be turned into first class quantum constraints. The conditions pointing to existence of anomalies were derived and it was analyzed how some kinds of anomalies can be removed. The conclusion was that the deformation quantization of pure Yang–Mills theory is straightforward whereas gravity is anomalous. It was also stated that in the ADM formalism of gravity, the anomaly is very complicated and the equations picking out physical states become infinite order functional differential equations. The Ashtekar variables remedy both of these problems in 2+1 dimensions but not in 3+1 dimensions. Recently, the deformation quantization was applied to linearized Einstein’s equations [22] using the analogy with Maxwell equations.

There is a series of works by C. Castro oriented to deformation quantization of gravity for strings and membranes contained in higher dimensions (in a sense, it is a realization of holographic idea): In Ref. [23], a geometric derivation of W_∞ gravity based on Fedosov’s deformation quantization of symplectic manifolds was elaborated. A holographic reduction of higher dimensional gravity was attained [24] based on the result of Ref. [25] that $m+n$ dimensional Einstein gravity can be identified with an m -dimensional generally invariant gauge theory of $Diff N$ (where N is an n -dimensional internal manifold), which allowed a deformation of gravity via Moyal non-commutative star products associated with the lower dimensional $SU(\infty)$ gauge theory. In [26] it was argued how a quantization of four dimensional gravity could be attained via a two dimensional quantum W_∞ gauge theory coupled to an infinite-component scalar-multiplet. It was shown also how strings and membranes actions in two and three dimensions emerge from four dimensional Einstein gravity by using the nonlinear connection formalism of Lagrange–Finsler and Hamilton–Cartan spaces (the formalism was also recently considered in quantum gravity in Refs. [1, 2, 3], see also the approaches with nonlinear connections to geometric mechanics [27, 28, 4] and applications in modern commutative and noncommutative gravity and string theories [30, 31]).

The present work belongs to a series of papers on Fedosov quantization of nonholonomic manifolds and generalized Lagrange–Finsler and Einstein spaces [1, 2, 3, 4, 29, 32]. Our main idea is to use such nonholonomic space-

time decompositions (for four dimensions, stating 2+2 splitting), with associated nonlinear connection (N-connection) induced by certain off-diagonal metric coefficients, when almost symplectic structures can be defined naturally for (pseudo) Riemannian spaces. Instead of the Levi-Civita connection, it is convenient to work with another class of metric-compatible linear connections adapted to the N-connection structure. Such connections are also uniquely defined by the metric structure but contain some nontrivial torsion components induced effectively by generic off-diagonal metric coefficients. The formalism allows us to quantize similarly both the Lagrange-Finsler and Einstein gravity. The final constructions can be redefined in terms of the Levi-Civita connection.

In Ref. [2], we proved that the Einstein gravity can be lifted canonically to the tangent bundle and transformed into an almost Kähler structure which can be quantized, for instance, following a generalization of Fedosov's method by Karabegov and Schlichenmaier [33]. It was emphasized that the models of quantum gravity on tangent bundles (or effectively constructed on higher dimension spacetimes) result in violation of local Lorentz symmetry. In addition to tangent bundles (extra dimensions) quantum deformation approaches to gravity, models with violated/restricted fundamental symmetries also present substantial interest in modern physics. In Ref. [3], we argued that Einstein's gravity can be quantized by nonholonomic deformations with effective generalized Lagrangians. This was done by transforming (semi) Riemannian metrics and connections into equivalent (almost symplectic) forms and connections adapted to a canonical N-connection structure on the same manifold, when the local Lorentz symmetry is preserved both for the classical and quantized models; see Ref. [3] for further details.

The aim of this work is twofold: First, to show how by nonholonomic frame transforms any (pseudo) Riemannian metric and corresponding Levi-Civita connection can be transformed into a canonical metric and nonlinear and linear connection structures for a Lagrange or Finsler geometry. We will also develop a technique for constructing an effective regular Lagrangian generated from a general metric structure with a formal 2+2 spacetime decomposition. Second, to prove that such effective Lagrange-Finsler spaces can be equivalently written in terms of an almost Kähler geometry. It is also analyzed how quantum properties of gravitational fields and motion equations are encoded into the 2-forms related the Chern-Weyl cohomological forms.

The work is organized as follows:

In section 2, we prove that any (pseudo) Riemannian metric can be represented in a form similar to canonical metrics in Lagrange or Finsler

geometry by the corresponding frame transform. We provide an analogous re-formulation of metric and frame variables from Einstein's gravity in Lagrange-Finsler type variables with an effective Lagrange or Finsler generating function defining a canonical 2+2 splitting of four dimensional spacetimes.

Section 3 is devoted to almost Kähler models of (pseudo) Riemannian and Lagrange spaces. Canonical almost symplectic forms and connections can be generated both by the metric structure and a nonholonomic distribution introduced on the original spacetime. We discuss the similarity and differences of fundamental geometric objects for the approaches with $3 + 1$ fibrations and $2 + 2$ nonholonomic splittings in gravity theories.

The Fedosov's operators are generalized in section 4 for nonholonomic manifolds and almost Kähler structures defined by effective Lagrange, or Finsler, fundamental functions on (pseudo) Riemannian manifolds.

In section 5, the main results on deformation quantization of almost Kähler geometries are reformulated for (pseudo) Riemannian nonholonomic manifolds and related Lagrange-Finsler spaces. We show how the information about Einstein's equations is encoded into the zero-degree cohomology coefficient of the correspondingly quantized Einstein's manifolds.

Finally, in section 6, we conclude and discuss the results.

2 (Pseudo)Riemannian Spaces Modeling Lagrange and Finsler Geometries

Let us consider a real (pseudo) Riemann manifold V^{2n} of necessary smooth class; $\dim V^{2n} = 2n$, where the dimension $n \geq 2$ is fixed.¹ We label the local coordinates in the form $u^\alpha = (x^i, y^a)$, or $u = (x, y)$, where indices run values $i, j, \dots = 1, 2, \dots, n$ and $a, b, \dots = n + 1, n + 2, \dots, n + n$, and x^i and y^a are respectively the conventional horizontal / holonomic (h) and vertical / nonholonomic coordinates (v). For the local Euclidean signature, we consider that all local basis vectors are real but, for the pseudo-Euclidean signature $(-, +, +, +)$, we introduce $e_{j=1} = i\partial/\partial x^1$, where i is the imaginary unity, $i^2 = -1$, and the local coordinate basis vectors can be written in the form $e_\alpha = \partial/\partial u^\alpha = (i\partial/\partial x^1, \partial/\partial x^2, \dots, \partial/\partial x^n, \partial/\partial y^a)$.² The Einstein's rule on summing up/low indices will be applied unless indicated otherwise.

¹for constructions related to Einstein's gravity $2n = 4$

²for simplicity, we shall omit to write in explicit form the imaginary unity considering that we can always distinguish the pseudo-Euclidean signature by a corresponding metric form or a local system of coordinates with a coordinate proportional to the imaginary unit

Any metric on V^{2n} can be written as

$$\mathbf{g} = g_{ij}(x, y) e^i \otimes e^j + h_{ab}(x, y) e^a \otimes e^b, \quad (1)$$

where the dual vielbeins (tetrads, in four dimensions) $e^a = (e^i, e^a)$ are parametrized as

$$e^i = e^i_{\underline{i}}(u) dx^{\underline{i}} \text{ and } e^a = e^a_{\underline{i}}(u) dx^{\underline{i}} + e^a_{\underline{a}}(u) dy^{\underline{a}}, \quad (2)$$

for $e_{\underline{\alpha}} = \partial/\partial u^{\underline{\alpha}} = (e_{\underline{i}} = \partial/\partial x^{\underline{i}}, e_{\underline{a}} = \partial/\partial y^{\underline{a}})$ and $e^{\underline{\beta}} = du^{\underline{\beta}} = (e^{\underline{j}} = dx^{\underline{j}}, dy^{\underline{b}})$ being, respectively, any fixed local coordinate base and dual base. One can also consider coordinate transforms such as $u^{\alpha'} = u^{\alpha'}(u^{\alpha}) = (x^{i'}(u^{\alpha}), y^{a'}(u^{\alpha}))$.

Proposition 2.1 *Any metric $\mathbf{g}$ (1) can be expressed in the form*

$$\check{\mathbf{g}} = \check{g}_{i'j'}(x, y) \check{e}^{i'} \otimes \check{e}^{j'} + \check{h}_{a'b'}(x, y) \check{\mathbf{e}}^{a'} \otimes \check{\mathbf{e}}^{b'}, \quad (3)$$

where $\check{e}^{i'} = \delta_{\underline{i}}^{i'} dx^{\underline{i}}$ and $\check{\mathbf{e}}^{a'} = \delta_{\underline{a}}^{a'}(u) dy^{\underline{a}} + \check{N}_{\underline{i}}^{a'}(u) dx^{\underline{i}}$ for

$$\check{h}_{a'b'}(u) = \frac{1}{2} \frac{\partial^2 \mathcal{L}(x^{i'}, y^{c'})}{\partial y^{a'} \partial y^{b'}}, \quad (4)$$

$$\check{N}_{\underline{i}}^{a'}(u) = \frac{\partial G^{a'}(x, y)}{\partial y^{n+\underline{i}}}, \quad (5)$$

where $\delta_{\underline{i}}^{i'}$ is the Kronecker symbol, $\check{g}_{i'j'} = \check{h}_{n+i' n+j'}$ and $\check{h}^{ab}$ is the inverse of $\check{h}_{a'b'}$, for $\det |\check{h}_{a'b'}| \neq 0$ and

$$2G^{a'}(x, y) = \frac{1}{2} \check{h}^{a' n+i} \left(\frac{\partial^2 \mathcal{L}}{\partial y^i \partial x^k} y^{n+k} - \frac{\partial \mathcal{L}}{\partial x^i} \right), \quad (6)$$

where $i, k = 1, 2, \dots, n$.

Proof. Let us fix a generating real function $\mathcal{L}(x^{i'}, y^{c'})$, called effective Lagrangian, and compute the values (4), (6) and (5), i.e., let's define the coefficients of metric (3) with respect to the local coordinate basis $du^{\underline{\alpha}} = (dx^{\underline{i}}, dy^{\underline{a}})$. Then we introduce the vielbein coefficients (2) in (1) and regroup the coefficients with respect to $du^{\underline{\alpha}} = (dx^{\underline{i}}, dy^{\underline{a}})$. Both formulas (1) and (3) define the same metric structure, i.e., $\mathbf{g} = \check{\mathbf{g}}$, if the following conditions are satisfied

$$g_{ij} e^i_{\underline{i}} e^j_{\underline{j}} = \check{g}_{\underline{i}\underline{j}}, \quad h_{ab} e^a_{\underline{a}} e^b_{\underline{b}} = \check{h}_{a'b'} \delta_{\underline{a}}^{a'} \delta_{\underline{b}}^{b'}, \quad h_{ab} e^a_{\underline{i}} e^b_{\underline{j}} = \check{h}_{a'b'} \check{N}_{\underline{i}}^{a'} \check{N}_{\underline{j}}^{b'}. \quad (7)$$

In four dimensions, $n = 2$, we have an algebraic system of 6 equations (7) for 12 unknown variables $e^i_{\underline{i}}, e^a_{\underline{a}}$ and $e^a_{\underline{a}}$, for $\check{g}_{i'j'}, \check{h}_{a'b'}$ and $\check{N}^{a'}_{\underline{a}}$ prescribed by $\mathcal{L}$. Such a system can be solved in order to find the 10 independent coefficients of any $\mathbf{g} = \{g_{\alpha\beta}\}$ (if this metric is a solution of the Einstein equations, there are only 6 independent coefficients, because of the Bianchi identities; four of the metric coefficients can be transformed to zero by use of the corresponding coordinate transforms). To build inverse constructions, we can prescribe the coefficients $g_{\alpha\beta}$ by taking any effective generating function $\mathcal{L}(x^{i'}, y^{c'})$ when the system (7) has nontrivial solutions (for simplicity, we can consider only real vielbeins and local coordinate bases adapted to the spacetime signature). $\square$

By a straightforward computation one proves:

Lemma 2.1 *Considering $\mathcal{L}$ from (4) and (6) to be a regular Lagrangian, we have that the Euler–Lagrange equations*

$$\frac{d}{d\tau} \left(\frac{\partial \mathcal{L}}{\partial y^i} \right) - \frac{\partial \mathcal{L}}{\partial x^i} = 0, \quad (8)$$

where $y^i = y^{n+i} = \frac{dx^i}{d\tau}$ for $x^i(\tau)$ depending on the parameter τ . The above Euler–Lagrange equations are equivalent to the “nonlinear” geodesic equations

$$\frac{d^2 x^i}{d\tau^2} + 2G^i(x^k, \frac{dx^j}{d\tau}) = 0 \quad (9)$$

defining the paths of a canonical semispray $S = y^i \frac{\partial}{\partial x^i} - 2G^a(x, y) \frac{\partial}{\partial y^a}$, for G^a given by equations (6).

The Lemma motivates:

Definition 2.1 *A (pseudo) Riemannian space with metric $\mathbf{g}$ (1) is modelled by a mechanical system with regular effective Lagrangian $\mathcal{L}$ if there is a nontrivial frame transform defined by any $e^i_{\underline{i}}, e^a_{\underline{a}}$ and $e^a_{\underline{a}}$ when $\mathbf{g} = \check{\mathbf{g}}$ (3).*

Inversely, we say that a regular mechanical model with Lagrangian $\mathcal{L}$ and Euler–Lagrange equations (8) is geometrized in terms of a (pseudo) Riemannian geometry with metric $\mathbf{g}$ (1) if $\mathcal{L}$ is a generating function for (4), (6) and (5), when $\mathbf{g} = \check{\mathbf{g}}$ (3) and the nonlinear geodesic equations (9) are equivalent to (8).

Any equivalent modelling of regular mechanical systems as (pseudo) Riemannian spaces introduces additional geometric structures on a manifold V^{2n} .

Definition 2.2 A nonlinear connection (N-connection) $\mathbf{N}$ on V^{2n} is defined by a Whitney sum (nonholonomic distribution)

$$T(V^{2n}) = h(V^{2n}) \oplus v(V^{2n}), \quad (10)$$

splitting globally the tangent bundle $T(V^{2n})$ into respective h - and v -subspaces, $h(V^{2n})$ and $v(V^{2n})$, given locally by a set of coefficients $N_i^a(x, y)$ where

$$\mathbf{N} = N_i^a(x, y) dx^i \otimes \frac{\partial}{\partial y^a}.$$

We note that a subclass of linear connections is defined by $N_i^a = \Gamma_b^a(x) y^b$.

Having prescribed on a $\mathbf{V}^{2n}$ a N-connection structure $\mathbf{N} = \{N_j^a\}$, we can define a preferred frame structure (with coefficients depending linearly on N_j^a) denoted $\mathbf{e}_\nu = (\mathbf{e}_i, e_a)$, where

$$\mathbf{e}_i = \frac{\partial}{\partial x^i} - N_i^a(u) \frac{\partial}{\partial y^a} \text{ and } e_a = \frac{\partial}{\partial y^a}, \quad (11)$$

with dual frame (coframe) structure $\mathbf{e}^\mu = (e^i, \mathbf{e}^a)$, where

$$e^i = dx^i \text{ and } \mathbf{e}^a = dy^a + N_i^a(u) dx^i, \quad (12)$$

satisfying nontrivial nonholonomy relations

$$[\mathbf{e}_\alpha, \mathbf{e}_\beta] = \mathbf{e}_\alpha \mathbf{e}_\beta - \mathbf{e}_\beta \mathbf{e}_\alpha = W_{\alpha\beta}^\gamma \mathbf{e}_\gamma \quad (13)$$

with (antisymmetric) anholonomy coefficients $W_{ia}^b = \partial_a N_i^b$ and $W_{ji}^a = \Omega_{ij}^a$.

Here boldface symbols are used for the spaces with N-connection structure and for the geometric objects adapted (N-adapted) to the h - and v -splitting (10). We can perform N-adapted geometric constructions by defining the coefficients of geometric objects (and associated equations) with respect to nonholonomic frames of type (11) and (12). The N-adapted tensors, vectors, forms, etc., are called respectively distinguished tensors, etc., (in brief, d-tensors, d-vectors, d-forms, etc.). For instance, a vector field $\mathbf{X} \in T\mathbf{V}^{2n}$ is expressed as $\mathbf{X} = (hX, vX)$, or $\mathbf{X} = X^\alpha \mathbf{e}_\alpha = X^i \mathbf{e}_i + X^a e_a$, where $hX = X^i \mathbf{e}_i$ and $vX = X^a e_a$ state, respectively, the horizontal (h) and vertical (v) components of the vector adapted to the N-connection structure.

Proposition 2.2 Any effective regular Lagrangian $\mathcal{L}$, prescribed on $\mathbf{V}^{2n}$, defines a canonical N-connection structure $\tilde{\mathbf{N}} = \{\tilde{N}_{\underline{i}}^{a'}(u)\}$ (5) and preferred frame structures $\tilde{\mathbf{e}}_\nu = (\tilde{\mathbf{e}}_i, e_{a'})$ and $\tilde{\mathbf{e}}^\mu = (e^i, \tilde{\mathbf{e}}^{a'})$.

Proof. The proposition can be proved by straightforward computations. The coefficients $\check{N}_{\underline{i}}^{a'}$ satisfy the conditions of Definition 2.2. We define $\check{\mathbf{e}}_\nu = (\check{\mathbf{e}}_i, e_a)$ and $\check{\mathbf{e}}^\mu = (e^i, \check{\mathbf{e}}^a)$ in explicit form by introducing $\check{N}_{\underline{i}}^{a'}$, respectively, in formulas (11) and (12). $\square$

Similar constructions can be defined for $\mathcal{L} = \mathcal{F}^2(x, y)$, where an effective Finsler metric $\mathcal{F}$ is a differentiable function of class C^∞ in any point (x, y) with $y \neq 0$ and is continuous in any point $(x, 0)$; $\mathcal{F}(x, y) > 0$ if $y \neq 0$; it satisfies the homogeneity condition $\mathcal{F}(x, \beta y) = |\beta| \mathcal{F}(x, y)$ for any nonzero $\beta \in \mathbb{R}$ and the Hessian (4) computed for $\mathcal{L} = \mathcal{F}^2$ is positive definite. In this case, we can say that a (pseudo) Riemannian space with metric $\mathbf{g}$ is modeled by an effective Finsler geometry and, inversely, a Finsler geometry is modeled on a (pseudo) Riemannian space. Such ideas were considered, for instance, in Ref. [28] for Lagrange and Finsler spaces defined on tangent bundles. In Ref. [31], we model such geometries on (pseudo) Riemannian and Riemann–Cartan spaces endowed with nonholonomic distributions.

Definition 2.3 *A (pseudo) Riemannian manifold $\mathbf{V}^{2n}$ is nonholonomic (N-anholonomic) if it is provided with a nonholonomic distribution on TV^{2n} (N-connection structure $\mathbf{N}$).*

We formulate the first main result in this paper:

Theorem 2.1 *Any (pseudo) Riemannian space can be transformed into a N-anholonomic manifold $\mathbf{V}^{2n}$ modeling an effective Lagrange (or Finsler) geometry by prescribing a generating Lagrange (or Finsler) function $\mathcal{L}(x, y)$ (or $\mathcal{F}(x, y)$).*

Proof. Such a proof follows from Propositions 2.1 and 2.2 and Lemma 2.1. It should be noted that, by corresponding vielbein transforms $e^i_{\underline{i}}, e^a_{\underline{a}}$ and $e^a_{\underline{a}}$, any metric $\mathbf{g}$ with coefficients defined with respect to an arbitrary co-frame $\mathbf{e}^\mu$, see (1), can be transformed into canonical Lagrange (Finsler) ones, $\check{\mathbf{g}}$ (3). The $\check{\mathbf{g}}$ coefficients are computed with respect to $\check{\mathbf{e}}^\mu = (e^i, \check{\mathbf{e}}^a)$, with the associated N-connection structure $\check{N}_{\underline{i}}^{a'}$, all defined by a prescribed $\mathcal{L}(x, y)$ (or $\mathcal{F}(x, y)$). $\square$

Finally, it should be noted that considering an arbitrary effective Lagrangian $\mathcal{L}(x, y)$ on a four dimensional (pseudo) Riemannian spacetime and defining a corresponding $2 + 2$ decomposition, local Lorentz invariance is not violated. We can work in any reference frame and coordinates, but the constructions adapted to the canonical N-connection structure and an analogous mechanical modeling are more convenient for developing a formalism

of deformation quantization of gravity following the appropriate methods for Lagrange–Finsler and almost Kähler spaces.

3 Almost Kähler Models for (Pseudo) Riemannian and Lagrange Spaces

The goal of this section is to prove that for any (pseudo) Riemannian metric and $n+n$ splitting we can define canonical almost symplectic structures. The analogous mechanical modeling developed in previous sections is important from two points of view: Firstly, it provides both geometric and physical interpretations for the class of nonholonomic transforms with $n+n$ splitting and adapting to the N–connection. Secondly, such canonical constructions can be equivalently redefined as a class of almost Kähler geometries with associated N–connection when certain symplectic forms and linear connection structures are canonically induced by the metric $\mathbf{g}(x, y)$ and effective Lagrangian $\mathcal{L}(x, y)$ on $\mathbf{V}^{2n}$.

3.1 Canonical Riemann–Lagrange symplectic structures

Let $\check{\mathbf{e}}_{\alpha'} = (\check{\mathbf{e}}_i, e_{b'})$ and $\check{\mathbf{e}}^{\alpha'} = (e^i, \check{\mathbf{e}}^{b'})$ be defined respectively by (11) and (12) for the canonical N–connection $\check{\mathbf{N}}$ stated by a metric structure $\mathbf{g} = \check{\mathbf{g}}$ on $\mathbf{V}^{2n}$. We introduce a linear operator $\check{\mathbf{J}}$ acting on tangent vectors to $\mathbf{V}^{2n}$ following formulas $\check{\mathbf{J}}(\check{\mathbf{e}}_i) = -e_{n+i}$ and $\check{\mathbf{J}}(e_{n+i}) = \check{\mathbf{e}}_i$, where the index α' runs values $n+i$ for $i = 1, 2, \dots, n$ and $\check{\mathbf{J}} \circ \check{\mathbf{J}} = -\mathbf{I}$ for $\mathbf{I}$ being the unity matrix. Equivalently, we introduce a tensor field on $\mathbf{V}^{2n}$,

$$\begin{aligned} \check{\mathbf{J}} &= \check{\mathbf{J}}^\alpha_\beta e_\alpha \otimes e^\beta = \check{\mathbf{J}}^\alpha_\beta \frac{\partial}{\partial u^\alpha} \otimes du^\beta \\ &= \check{\mathbf{J}}^{\alpha'}_{\beta'} \check{\mathbf{e}}_{\alpha'} \otimes \check{\mathbf{e}}^{\beta'} = -e_{n+i} \otimes e^i + \check{\mathbf{e}}_i \otimes \check{\mathbf{e}}^{n+i} \\ &= -\frac{\partial}{\partial y^i} \otimes dx^i + \left(\frac{\partial}{\partial x^i} - \check{N}_i^{n+j} \frac{\partial}{\partial y^j} \right) \otimes \left(dy^i + \check{N}_k^{n+i} dx^k \right). \end{aligned}$$

It is clear that $\check{\mathbf{J}}$ defines globally an almost complex structure on $\mathbf{V}^{2n}$ completely determined by a fixed $\mathcal{L}(x, y)$. Using vielbeins $\mathbf{e}^\alpha_{\underline{\alpha}}$ and their duals $\mathbf{e}^{\underline{\alpha}}_\alpha$, defined by $e^i_{\underline{i}}, e^a_{\underline{a}}$ and $e^a_{\underline{a}}$ as a solution of (7), we can compute the coefficients $\check{\mathbf{J}}$ with respect to any local basis e_α and e^α on $\mathbf{V}^{n+n}$, $\check{\mathbf{J}}^\alpha_\beta = \mathbf{e}^\alpha_{\underline{\alpha}} \check{\mathbf{J}}^\alpha_{\underline{\beta}} \mathbf{e}^{\underline{\beta}}_\beta$. In general, we can define an almost complex structure $\mathbf{J}$ for an arbitrary N–connection $\mathbf{N}$ by using N–adapted bases (11) and (12), not necessarily induced by an effective Lagrange function.

Definition 3.1 *The Nijenhuis tensor field for any almost complex structure $\mathbf{J}$ determined by a N -connection (equivalently, the curvature of N -connection) is defined as*

$$\mathbf{J}\Omega(\mathbf{X}, \mathbf{Y}) = -[\mathbf{X}, \mathbf{Y}] + [\mathbf{JX}, \mathbf{JY}] - \mathbf{J}[\mathbf{JX}, \mathbf{Y}] - \mathbf{J}[\mathbf{X}, \mathbf{JY}], \quad (14)$$

for any d -vectors $\mathbf{X}$ and $\mathbf{Y}$.

With respect to N -adapted bases (11) and (12) the Neijenhuis tensor $\mathbf{J}\Omega = \{\Omega_{ij}^a\}$ has the coefficients

$$\Omega_{ij}^a = \frac{\partial N_i^a}{\partial x^j} - \frac{\partial N_j^a}{\partial x^i} + N_i^b \frac{\partial N_j^a}{\partial y^b} - N_j^b \frac{\partial N_i^a}{\partial y^b}. \quad (15)$$

A N -anholonomic manifold $\mathbf{V}^{2n}$ is integrable if $\Omega_{ij}^a = 0$. We get a complex structure if and only if both the h - and v -distributions are integrable, i.e., if and only if $\Omega_{ij}^a = 0$ and $\frac{\partial N_j^a}{\partial y^i} - \frac{\partial N_i^a}{\partial y^j} = 0$.

Definition 3.2 *An almost symplectic structure on a manifold V^{n+m} , $\dim V^{n+m} = n+m$, is defined by a nondegenerate 2-form $\theta = \frac{1}{2}\theta_{\alpha\beta}(u)e^\alpha \wedge e^\beta$.*

We have

Proposition 3.1 *For any θ on V^{n+m} , there is a unique N -connection $\mathbf{N} = \{N_i^a\}$ defined as a splitting $TV^{n+m} = hV^{n+m} \oplus vV^{n+m}$, where indices $i, j, \dots = 1, 2, \dots, n$ and $a, b, \dots = n+1, n+1, \dots, n+m$. The function θ satisfies the following conditions:*

$$\theta = (h\mathbf{X}, v\mathbf{Y}) = 0 \text{ and } \theta = h\theta + v\theta, \quad (16)$$

for any $\mathbf{X} = h\mathbf{X} + v\mathbf{X}$, $\mathbf{Y} = h\mathbf{Y} + v\mathbf{Y}$ and $h\theta(\mathbf{X}, \mathbf{Y}) \doteq \theta(h\mathbf{X}, h\mathbf{Y})$, $v\theta(\mathbf{X}, \mathbf{Y}) \doteq \theta(v\mathbf{X}, v\mathbf{Y})$. Here the symbol " $\doteq$ " means "by definition".

Proof. For $\mathbf{X} = \mathbf{e}_\alpha = (\mathbf{e}_i, e_a)$ and $\mathbf{Y} = \mathbf{e}_\beta = (\mathbf{e}_l, e_b)$, where $\mathbf{e}_\alpha$ is a N -adapted basis of type (11) of dimension $n+m$, we write the first equation in (16) as $\theta = \theta(\mathbf{e}_i, e_a) = \theta(\frac{\partial}{\partial x^i}, \frac{\partial}{\partial y^a}) - N_i^b \theta(\frac{\partial}{\partial y^b}, \frac{\partial}{\partial y^a}) = 0$. We can find a unique solution form and define N_i^b if $\text{rank}|\theta(\frac{\partial}{\partial y^b}, \frac{\partial}{\partial y^a})| = m$. Denoting locally

$$\theta = \frac{1}{2}\theta_{ij}(u)e^i \wedge e^j + \frac{1}{2}\theta_{ab}(u)\mathbf{e}^a \wedge \mathbf{e}^b, \quad (17)$$

where the first term is for $h\theta$ and the second term is $v\theta$, we get the second formula in (16). We may consider the particular case in which $n = m$. $\square$

Definition 3.3 An almost Hermitian model of a (pseudo) Riemannian space V^{2n} equipped with an N -connection structure $\mathbf{N}$ is defined by a triple $\mathbf{H}^{2n} = (V^{2n}, \theta, \mathbf{J})$, where $\theta(\mathbf{X}, \mathbf{Y}) \doteq \mathbf{g}(\mathbf{J}\mathbf{X}, \mathbf{Y})$.

In addition, we have

Definition 3.4 A space $\mathbf{H}^{2n}$ is almost Kähler, denoted $\mathbf{K}^{2n}$, if and only if $d\theta = 0$.

If a (pseudo) Riemannian space is modeled by a Lagrange–Finsler geometry, the second main result of this paper follows

Theorem 3.1 Having chosen a generating function $\mathcal{L}(x, y)$ (or $\mathcal{F}(x, y)$) on a (pseudo) Riemannian manifold V^{n+n} , we can model this space as an almost Kähler geometry, i.e. $\check{\mathbf{H}}^{2n} = \check{\mathbf{K}}^{2n}$.

Proof. For $\mathbf{g} = \check{\mathbf{g}}$ (3) and structures $\check{\mathbf{N}}$ and $\check{\mathbf{J}}$ canonically defined by $\mathcal{L}$, we define $\check{\theta}(\mathbf{X}, \mathbf{Y}) \doteq \check{\mathbf{J}}(\check{\mathbf{F}}\mathbf{X}, \mathbf{Y})$ for any d-vectors $\mathbf{X}$ and $\mathbf{Y}$. In local N -adapted form form, we have

$$\begin{aligned} \check{\theta} &= \frac{1}{2} \check{\theta}_{\alpha\beta}(u) e^\alpha \wedge e^\beta = \frac{1}{2} \check{\theta}_{\underline{\alpha}\underline{\beta}}(u) du^\alpha \wedge du^\beta \\ &= \check{g}_{ij}(x, y) \check{e}^{n+i} \wedge dx^j = \check{g}_{ij}(x, y) (dy^{n+i} + \check{N}_k^{n+i} dx^k) \wedge dx^j. \end{aligned} \quad (18)$$

Let us consider the form $\check{\omega} = \frac{1}{2} \frac{\partial \mathcal{L}}{\partial y^{n+i}} dx^i$. A straightforward computation, using Proposition 2.2 and N -connection $\check{\mathbf{N}}$ (5), shows that $\check{\theta} = d\check{\omega}$, which means that $d\check{\theta} = dd\check{\omega} = 0$ and that the canonical effective Lagrange structures $\mathbf{g} = \check{\mathbf{g}}$, $\check{\mathbf{N}}$ and $\check{\mathbf{J}}$ induce an almost Kähler geometry. Instead of "Lagrangian mechanics variables" we can introduce another type redefining $\check{\theta}$ with respect to an arbitrary co-frame basis using vielbeins $\mathbf{e}^\alpha_{\underline{\alpha}}$ and their duals $\mathbf{e}^{\underline{\alpha}}_\alpha$, defined by $e^i_{\underline{i}}, e^a_{\underline{a}}$ and $e^a_{\underline{a}}$ (7). So, we can compute $\check{\theta}_{\alpha\beta} = \mathbf{e}^\alpha_{\underline{\alpha}} \mathbf{e}^\beta_{\underline{\beta}} \check{\theta}_{\underline{\alpha}\underline{\beta}}$ and express the 2-form (18) as $\check{\theta} = \frac{1}{2} \check{\theta}_{ij}(u) e^i \wedge e^j + \frac{1}{2} \check{\theta}_{ab}(u) \check{\mathbf{e}}^a \wedge \check{\mathbf{e}}^b$, see (17). The coefficients $\check{\theta}_{ab} = \check{\theta}_{n+i \ n+j}$ above are equal, respectively, to the coefficients $\check{\theta}_{ij}$ and the dual N -adapted basis $\check{\mathbf{e}}^\alpha = (e^i, \check{\mathbf{e}}^a)$ is elongated by $\check{N}_j^a$ (5). It should be noted that for a general 2-form θ directly constructed from a metric $\mathbf{g}$ and almost complex $\mathbf{J}$ structures on V^{2n} , we have that $d\theta \neq 0$. For a $n+n$ splitting induced by an effective Lagrange (Finsler) generating function, we have $d\check{\theta} = 0$ which results in a canonical almost Kähler model completely defined by $\mathbf{g} = \check{\mathbf{g}}$ and chosen $\mathcal{L}(x, y)$ (or $\mathcal{F}(x, y)$). $\square$

3.2 N-adapted symplectic connections

In our approach, we work with nonholonomic (pseudo) Riemannian manifolds $\mathbf{V}^{2n}$ enabled with an effective N-connection and almost symplectic structures defined canonically by the metric structure $\mathbf{g} = \check{\mathbf{g}}$ and a fixed $\mathcal{L}(x, y)$. In this section, we analyze the class of linear connections that can be adapted to the N-connection and/or symplectic structure and defined canonically if a corresponding nonholonomic distribution is induced completely by $\mathcal{L}$, or $\mathcal{F}$.

From the class of arbitrary affine connections on $\mathbf{V}^{2n}$, one prefers to work with N-adapted linear connections, called distinguished connections (d-connections).

Definition 3.5 *A linear connection on $\mathbf{V}^{2n}$ is a d-connection*

$$\mathbf{D} = (hD; vD) = \{\Gamma_{\beta\gamma}^\alpha = (L_{jk}^i, {}^vL_{bk}^a; C_{jc}^i, {}^vC_{bc}^a)\},$$

with local coefficients computed with respect to (11) and (12), which preserves the distribution (10) under parallel transports.

For a d-connection $\mathbf{D}$, we can define respectively the torsion and curvature tensors,

$$\mathbf{T}(\mathbf{X}, \mathbf{Y}) \doteq \mathbf{D}_\mathbf{X}\mathbf{Y} - \mathbf{D}_\mathbf{Y}\mathbf{X} - [\mathbf{X}, \mathbf{Y}], \quad (19)$$

$$\mathbf{R}(\mathbf{X}, \mathbf{Y})\mathbf{Z} \doteq \mathbf{D}_\mathbf{X}\mathbf{D}_\mathbf{Y}\mathbf{Z} - \mathbf{D}_\mathbf{Y}\mathbf{D}_\mathbf{X}\mathbf{Z} - \mathbf{D}_{[\mathbf{X}, \mathbf{Y}]}\mathbf{Z}, \quad (20)$$

where $[\mathbf{X}, \mathbf{Y}] \doteq \mathbf{X}\mathbf{Y} - \mathbf{Y}\mathbf{X}$, for any d-vectors $\mathbf{X}$ and $\mathbf{Y}$. The coefficients $\mathbf{T} = \{\mathbf{T}_{\beta\gamma}^\alpha\}$ and $\mathbf{R} = \{\mathbf{R}_{\beta\gamma\tau}^\alpha\}$ can be written in terms of $\mathbf{e}_\alpha$ and $\mathbf{e}^\beta$ by introducing $\mathbf{X} \rightarrow \mathbf{e}_\alpha$, $\mathbf{Y} \rightarrow \mathbf{e}_\beta$, $\mathbf{Z} \rightarrow \mathbf{e}_\gamma$ in (19) and (20), see Ref. [31] for details.

Definition 3.6 *A d-connection $\mathbf{D}$ is metric compatible with a d-metric $\mathbf{g}$ if $\mathbf{D}_\mathbf{X}\mathbf{g} = 0$ for any d-vector field $\mathbf{X}$.*

If an almost symplectic structure is defined on a N-anholonomic manifold, one considers:

Definition 3.7 *An almost symplectic d-connection ${}_\theta\mathbf{D}$ on $\mathbf{V}^{2n}$, or (equivalently) a d-connection compatible with an almost symplectic structure θ , is defined such that ${}_\theta\mathbf{D}$ is N-adapted, i.e., it is a d-connection, and ${}_\theta\mathbf{D}_\mathbf{X}\theta = 0$, for any d-vector $\mathbf{X}$.*

We can always fix a d-connection ${}_{\circ}\mathbf{D}$ on $\mathbf{V}^{2n}$ and then construct an almost symplectic ${}_{\theta}\mathbf{D}$.

Example 3.1 *Let us represent θ in N-adapted form (17). Having chosen a*

$$\begin{aligned} {}_{\circ}\mathbf{D} &= \{h \circ D = ({}_{\circ}D_k, {}^{\circ}D_k); v \circ D = ({}_{\circ}D_c, {}^{\circ}D_c)\} \\ &= \{ {}_{\circ}\mathbf{\Gamma}_{\beta\gamma}^{\alpha} = ({}_{\circ}L_{jk}^i, {}^{\circ}L_{bk}^a; {}_{\circ}C_{jc}^i, {}^{\circ}C_{bc}^a) \}, \end{aligned}$$

we can verify that

$$\begin{aligned} {}_{\theta}\mathbf{D} &= \{h {}_{\theta}D = ({}_{\theta}D_k, {}^{\theta}D_k); v {}_{\theta}D = ({}_{\theta}D_c, {}^{\theta}D_c)\} \\ &= \{ {}_{\theta}\mathbf{\Gamma}_{\beta\gamma}^{\alpha} = ({}_{\theta}L_{jk}^i, {}^{\theta}L_{bk}^a; {}_{\theta}C_{jc}^i, {}^{\theta}C_{bc}^a) \}, \end{aligned}$$

with

$$\begin{aligned} {}_{\theta}L_{jk}^i &= {}_{\circ}L_{jk}^i + \frac{1}{2}\theta^{ih} {}_{\circ}D_k\theta_{jh}, \quad {}^{\theta}L_{bk}^a = {}^{\circ}L_{bk}^a + \frac{1}{2}\theta^{ae} {}^{\circ}D_k\theta_{eb}, \\ {}_{\theta}C_{jc}^i &= {}_{\circ}C_{jc}^i + \frac{1}{2}\theta^{ih} {}_{\circ}D_c\theta_{jh}, \quad {}^{\theta}C_{bc}^a = {}^{\circ}C_{bc}^a + \frac{1}{2}\theta^{ae} {}^{\circ}D_c\theta_{eb}, \end{aligned} \quad (21)$$

satisfies the conditions ${}_{\theta}D_k\theta_{jh} = 0$, ${}^{\theta}D_k\theta_{eb} = 0$, ${}_{\theta}D_c\theta_{jh} = 0$, ${}^{\theta}D_c\theta_{eb} = 0$, which is equivalent to ${}_{\theta}\mathbf{D}\mathbf{X}\theta = 0$ from Definition 3.7.

Let us introduce the operators

$$\Theta_{jk}^{hi} = \frac{1}{2}(\delta_j^h \delta_k^i - \theta_{jk}\theta^{ih}) \text{ and } \Theta_{cd}^{ab} = \frac{1}{2}(\delta_c^a \delta_d^b - \theta_{cd}\theta^{ab}), \quad (22)$$

with the coefficients computed with respect to N-adapted bases (11) and (12). By straightforward computations, one proves the following theorem.

Theorem 3.2 *The set of d-connections ${}_s\mathbf{\Gamma}_{\beta\gamma}^{\alpha} = ({}_sL_{jk}^i, {}^sL_{bk}^a; {}_sC_{jc}^i, {}^sC_{bc}^a)$, compatible with an almost symplectic structure θ (17), are parametrized by*

$$\begin{aligned} {}_sL_{jk}^i &= {}_{\theta}L_{jk}^i + \Theta_{jl}^{hi} Y_{hk}^l, \quad {}^sL_{bk}^a = {}^{\theta}L_{bk}^a + \Theta_{bd}^{ca} Y_{ck}^d, \\ {}_sC_{jc}^i &= {}_{\theta}C_{jc}^i + \Theta_{jl}^{hi} Y_{hc}^l, \quad {}^sC_{bc}^a = {}^{\theta}C_{bc}^a + \Theta_{bd}^{ea} Y_{ec}^d, \end{aligned} \quad (23)$$

where ${}_{\theta}\mathbf{\Gamma}_{\beta\gamma}^{\alpha} = ({}_{\theta}L_{jk}^i, {}^{\theta}L_{bk}^a; {}_{\theta}C_{jc}^i, {}^{\theta}C_{bc}^a)$ is given by (21), the Θ -operators are those from (22) and $\mathbf{Y}_{\beta\gamma}^{\alpha} = (Y_{jk}^i, Y_{bk}^a, Y_{jc}^i, Y_{bc}^a)$ are arbitrary d-tensor fields.

From the set of metric and/or almost symplectic compatible d-connections on a (pseudo) Riemannian manifold V^{2n} , we can select those which are completely defined by $\mathbf{g}$ and a prescribed effective Lagrange structure $\mathcal{L}(x, y)$:

Theorem 3.3 *There is a unique normal d-connection*

$$\begin{aligned}\widehat{\mathbf{D}} &= \left\{ h\widehat{D} = (\widehat{D}_k, {}^v\widehat{D}_k = \widehat{D}_k); {}^v\widehat{D} = (\widehat{D}_c, {}^v\widehat{D}_c = \widehat{D}_c) \right\} \\ &= \left\{ \widehat{\Gamma}_{\beta\gamma}^\alpha = (\widehat{L}_{jk}^i, {}^v\widehat{L}_{n+j \ n+k}^{n+i} = \widehat{L}_{jk}^i; \widehat{C}_{jc}^i = {}^v\widehat{C}_{n+j \ c}^{n+i}, {}^v\widehat{C}_{bc}^a = \widehat{C}_{bc}^a) \right\},\end{aligned}\quad (24)$$

which is metric compatible, $\widehat{D}_k\check{g}_{ij} = 0$ and $\widehat{D}_c\check{g}_{ij} = 0$, and completely defined by $\mathbf{g} = \check{\mathbf{g}}$ and a fixed $\mathcal{L}(x, y)$.

Proof. First, we note that if a normal d-connection exists, it is completely defined by couples of h- and v-components $\widehat{\mathbf{D}}_\alpha = (\widehat{D}_k, \widehat{D}_c)$, i.e. $\widehat{\Gamma}_{\beta\gamma}^\alpha = (\widehat{L}_{jk}^i, {}^v\widehat{C}_{bc}^a)$. Choosing

$$\widehat{L}_{jk}^i = \frac{1}{2}\check{g}^{ih}(\check{\mathbf{e}}_k\check{g}_{jh} + \check{\mathbf{e}}_j\check{g}_{hk} - \check{\mathbf{e}}_h\check{g}_{jk}), \widehat{C}_{jk}^i = \frac{1}{2}\check{g}^{ih}\left(\frac{\partial\check{g}_{jh}}{\partial y^k} + \frac{\partial\check{g}_{hk}}{\partial y^j} - \frac{\partial\check{g}_{jk}}{\partial y^h}\right), \quad (25)$$

where $\check{\mathbf{e}}_k = \partial/\partial x^k + \check{N}_k^a\partial/\partial y^a$, $\check{N}_k^a$ and $\check{g}_{jk} = \check{h}_{n+i \ n+j}$ are defined by canonical values (4) and (5) induced by a regular $\mathcal{L}(x, y)$, we can prove that this d-connection is unique and satisfies the conditions of the theorem. Using vielbeins $\mathbf{e}_{\underline{\alpha}}^\alpha$ and their duals $\mathbf{e}_{\underline{\alpha}}^\alpha$, defined by $e_{\underline{i}}^i, e_{\underline{j}}^j$ and $e_{\underline{a}}^a$ satisfying (7), we can compute the coefficients of $\widehat{\Gamma}_{\beta\gamma}^\alpha$ (24) with respect to arbitrary frame basis e_α and co-basis e^α on V^{n+m} . $\square$

Introducing the normal d-connection 1-form $\widehat{\Gamma}_j^i = \widehat{L}_{jk}^i e^k + \widehat{C}_{jk}^i \check{\mathbf{e}}^k$, for $e^k = dx^k$ and $\check{\mathbf{e}}^k = dy^k + \check{N}_i^k dx^i$, we can prove that the Cartan structure equations are satisfied,

$$de^k - e^j \wedge \widehat{\Gamma}_j^k = -\widehat{\mathcal{T}}^i, \quad d\check{\mathbf{e}}^k - \check{\mathbf{e}}^j \wedge \widehat{\Gamma}_j^k = -{}^v\widehat{\mathcal{T}}^i, \quad (26)$$

and

$$d\widehat{\Gamma}_j^i - \widehat{\Gamma}_j^h \wedge \widehat{\Gamma}_h^i = -\widehat{\mathcal{R}}^i_j. \quad (27)$$

The h- and v-components of the torsion 2-form $\widehat{\mathcal{T}}^\alpha = (\widehat{\mathcal{T}}^i, {}^v\widehat{\mathcal{T}}^i) = \widehat{\mathbf{T}}_{\tau\beta}^\alpha \check{\mathbf{e}}^\tau \wedge \check{\mathbf{e}}^\beta$ and from (26) the components are computed

$$\widehat{\mathcal{T}}^i = \widehat{C}_{jk}^i e^j \wedge \check{\mathbf{e}}^k, \quad {}^v\widehat{\mathcal{T}}^i = \frac{1}{2}\check{\Omega}_{kj}^i e^k \wedge e^j + \left(\frac{\partial\check{N}_k^i}{\partial y^j} - \widehat{L}_{kj}^i\right) e^k \wedge \check{\mathbf{e}}^j, \quad (28)$$

where $\check{\Omega}_{kj}^i$ are coefficients of the curvature of the canonical N-connection $\check{N}_k^i$ defined by formulas similar to (15). Such formulas also follow from (19) redefined for $\widehat{\mathbf{D}}_\alpha$ and $\check{\mathbf{e}}_\alpha$, when the torsion $\widehat{\mathbf{T}}_{\beta\gamma}^\alpha$ is parametrized as

$$\widehat{T}_{jk}^i = 0, \widehat{T}_{jc}^i = \widehat{C}_{jc}^i, \widehat{T}_{ij}^a = \check{\Omega}_{ij}^a, \widehat{T}_{ib}^a = e_b \check{N}_i^a - \widehat{L}_{bi}^a, \widehat{T}_{bc}^a = 0. \quad (29)$$

It should be noted that $\hat{\mathbf{T}}$ vanishes on h- and v-subspaces, i.e. $\hat{T}_{jk}^i = 0$ and $\hat{T}_{bc}^a = 0$, but certain nontrivial h-v-components induced by the nonholonomic structure are defined canonically by $\mathbf{g} = \check{\mathbf{g}}$ and $\mathcal{L}$.

We can also compute the curvature 2-form from (27),

$$\hat{\mathcal{R}}^\tau_\gamma = \hat{\mathbf{R}}^\tau_{\gamma\alpha\beta} \check{\mathbf{e}}^\alpha \wedge \check{\mathbf{e}}^\beta = \frac{1}{2} \hat{R}^i_{jkh} e^k \wedge e^h + \hat{P}^i_{jka} e^k \wedge \check{\mathbf{e}}^a + \frac{1}{2} \hat{S}^i_{jcd} \check{\mathbf{e}}^c \wedge \check{\mathbf{e}}^d, \quad (30)$$

where the nontrivial N-adapted coefficients of curvature $\hat{\mathbf{R}}^\alpha_{\beta\gamma\tau}$ of $\hat{\mathbf{D}}$ are (such formulas can be proven also from (20) written for $\hat{\mathbf{D}}_\alpha$ and $\check{\mathbf{e}}_\alpha$)

$$\begin{aligned} \hat{R}^i_{hjk} &= \check{\mathbf{e}}_k \hat{L}^i_{hj} - \check{\mathbf{e}}_j \hat{L}^i_{hk} + \hat{L}^m_{hj} \hat{L}^i_{mk} - \hat{L}^m_{hk} \hat{L}^i_{mj} - \hat{C}^i_{ha} \check{\Omega}^a_{kj}, \\ \hat{P}^i_{jka} &= e_a \hat{L}^i_{jk} - \hat{\mathbf{D}}_k \hat{C}^i_{ja}, \quad \hat{S}^a_{bcd} = e_d \hat{C}^a_{bc} - e_c \hat{C}^a_{bd} + \hat{C}^e_{bc} \hat{C}^a_{ed} - \hat{C}^e_{bd} \hat{C}^a_{ec}. \end{aligned} \quad (31)$$

If instead of an effective Lagrange function one considers a Finsler generating fundamental function $\mathcal{F}^2$, similar formulas for the torsion and curvature of the normal d-connection can also be found.

There is another very important property of the normal d-connection:

Theorem 3.4 *The normal d-connection $\hat{\mathbf{D}}$ defines a unique almost symplectic d-connection, $\hat{\mathbf{D}} \equiv {}_\theta \hat{\mathbf{D}}$, see Definition 3.7, which is N-adapted, i.e. it preserves under parallelism the splitting (10), ${}_\theta \hat{\mathbf{D}}_{\mathbf{X}} \check{\theta} = 0$ and $\hat{T}_{jk}^i = \hat{T}_{bc}^a = 0$, i.e. the torsion is of type (29).*

Proof. Applying the conditions of the theorem to the coefficients (25), the proof follows in a straightforward manner. $\square$

It is pertinent to note that the normal d-connection $\hat{\Gamma}^\alpha_{\beta\gamma}$ (24) is a N-anholonomic analog of the affine connection ${}^K \Gamma^\alpha_{\beta\gamma}$ and Nijenhuis tensor ${}^K \Omega^\alpha_{\beta\gamma}$ with the torsion satisfying the condition ${}^K \mathbf{T}^\alpha_{\beta\gamma} = (1/4) {}^K \Omega^\alpha_{\beta\gamma}$, as considered in Ref. [33]. For trivial N-connection structures, by corresponding frame and coordinate transforms, we can identify $\hat{\Gamma}^\alpha_{\beta\gamma}$ with ${}^K \Gamma^\alpha_{\beta\gamma}$ (we used this property in our former works [1, 2, 3, 4]).

In this section, we proved that a N-adapted and almost symplectic $\hat{\Gamma}^\alpha_{\beta\gamma}$ can be uniquely defined by a (pseudo) Riemannian metric $\mathbf{g}$ if we prescribe an effective Lagrange, or Finsler, function $\mathcal{L}$, or $\mathcal{F}$ on V^{2n} . This allows us to construct an analogous Lagrange model for gravity and, at the next step, to transform it equivalently in an almost Kähler structure adapted to a corresponding $n + n$ spacetime splitting. For the Einstein metrics, we get a canonical $2 + 2$ decomposition for which we can apply the Fedosov's quantization if the geometric objects and operators are adapted to the associated N-connection.

Definition 3.8 *A (pseudo) Riemannian space is described in Lagrange–Finsler variables if its vielbein, metric and linear connection structures are equivalently transformed into corresponding canonical N –connection, Lagrange–Finsler metric and normal / almost symplectic d –connection structures.*

It should be noted that former approaches to the canonical and quantum loop quantization of gravity were elaborated for $3 + 1$ fibrations and corresponding ADM and Ashtekar variables with further modifications. On the other hand, in order to elaborate certain approaches to deformation quantization of gravity, it is crucial to work with nonholonomic $2 + 2$ structures, which is more convenient for certain Lagrange geometrized constructions and their almost symplectic variants. For other models, the $3 + 1$ splitting preserves a number of similarities to Hamilton mechanics. In our approach, the spacetime decompositions are defined by corresponding N –connection structures, which can be induced canonically by effective Lagrange, or Finsler, generating functions. One works both with N –adapted metric coefficients and nonholonomic frame coefficients, the last ones being defined by generic off–diagonal metric coefficients and related N –connection coefficients. In the models related to $3 + 1$ fibrations, one works with shift functions and frame variables which contain all dynamical information, instead of metrics.

We also discuss here the similarities and differences of preferred classes of linear connections used for $3 + 1$ and $2 + 2$ structures. In the first case, the Ashtekar variables (and further modifications) were proved to simplify the constraint structure of a gauge like theory to which the Einstein theory was transformed in order to develop a background independent quantization of gravity. In the second case, the analogs of Ashtekar variables are generated by a canonical Lagrange–Finsler type metric and/or corresponding almost symplectic structure, both adapted to the N –connection structure. It is also involved the normal d –connection which is compatible with the almost symplectic structure and completely defined by the metric structure, alternatively to the Levi–Civita connection (the last one is not adapted to the N –connection and induced almost symplectic structure). In fact, all constructions for the normal d –connection can be redefined in an equivalent form to the Levi–Civita connection (see below section 5.2 and Refs. [30, 31]), or in Ashtekar variables, but in such cases the canonical $2 + 2$ splitting and almost Kähler structure are mixed by general frame and linear connection deformations.

Finally, it should be noted that in our approach we are inspired by a number of results and methods from Finsler and Lagrange geometry. For

instance, the original proofs that Finsler and Lagrange geometries are equivalent to certain classes of almost Kähler geometries with N-connection structures were obtained in Refs. [34, 35], see also reviews [27, 28, 31]. But those constructions were elaborated for tangent bundles which are not related to standard models of modern physics. Re-defining the constructions for nonholonomic structures on classical and quantum spacetime models, we could develop new important and effective methods from the geometry of nonholonomic manifolds and apply them to deformation quantization of gravity.

4 Distinguished Fedosov's Operators

The Fedosov's approach to deformation quantization [5, 6] will be extended for (pseudo) Riemannian manifolds V^{2n} endowed with an effective Lagrange function $\mathcal{L}$. The constructions elaborated in Ref. [33] will be adapted to the canonical N-connection structure by considering decompositions with respect to $\check{\mathbf{e}}_\nu = (\check{\mathbf{e}}_i, e_{a'})$ and $\check{\mathbf{e}}^\mu = (e^i, \check{\mathbf{e}}^{a'})$ defined by a metric $\mathbf{g}$ (1). For simplicity, we shall work only with the normal/ almost symplectic d-connection, $\hat{\mathbf{D}} \equiv {}_\theta \hat{\mathbf{D}}$ (24), see Definition 3.7, but it should be emphasized here that we can use any d-connection from the family (23) and develop a corresponding deformation quantization. Usually, the proofs referring to constructions not adapted to N-connections [33], and on Lagrange (Finsler) spaces related to quantum gravity models on tangent bundles [2] will be sketched, while the details can be found in the corresponding references. In this work, the formulas are redefined on nonholonomic (pseudo) Riemannian manifolds modeling effective regular mechanical systems and corresponding almost Kähler structures.

We introduce the tensor $\check{\mathbf{A}}^{\alpha\beta} \doteq \check{\theta}^{\alpha\beta} - i \check{\mathbf{g}}^{\alpha\beta}$, where $\check{\theta}^{\alpha\beta}$ is the form (18) with "up" indices and $\check{\mathbf{g}}^{\alpha\beta}$ is the inverse to $\check{\mathbf{g}}_{\alpha\beta}$ stated by coefficients of (3). The local coordinates on $\mathbf{V}^{2n}$ are parametrized as $u = \{u^\alpha\}$ and the local coordinates on $T_u \mathbf{V}^{2n}$ are labeled $(u, z) = (u^\alpha, z^\beta)$, where z^β are fiber coordinates.

The formalism of deformation quantization can be developed by using $C^\infty(V)[[v]]$, the space of formal series of variable v with coefficients from $C^\infty(V)$ on a Poisson manifold $(V, \{\cdot, \cdot\})$ (in this work, we deal with an almost Poisson structure defined by the canonical almost symplectic structure). One defines an associative algebra structure on $C^\infty(V)[[v]]$ with a v -linear and v -adically continuous star product

$${}^1f * {}^2f = \sum_{r=0}^{\infty} {}_r C({}^1f, {}^2f) v^r, \quad (32)$$

where ${}_r C, r \geq 0$, are bilinear operators on $C^\infty(V)$ with ${}_0 C({}^1 f, {}^2 f) = {}^1 f {}^2 f$ and ${}_1 C({}^1 f, {}^2 f) - {}_1 C({}^2 f, {}^1 f) = i\{{}^1 f, {}^2 f\}$; i being the complex unity. Constructions of type (32) are used for stating a formal Wick product

$$a \circ b(z) \doteq \exp \left(i \frac{v}{2} \tilde{\Lambda}^{\alpha\beta} \frac{\partial^2}{\partial z^\alpha \partial z_{[1]}^\beta} \right) a(z) b(z_{[1]})|_{z=z_{[1]}}, \quad (33)$$

for two elements a and b defined by series of type

$$a(v, z) = \sum_{r \geq 0, |\{\alpha\}| \geq 0} a_{r, \{\alpha\}}(u) z^{\{\alpha\}} v^r, \quad (34)$$

where by $\{\alpha\}$ we label a multi-index. This way, we define a formal Wick algebra $\tilde{\mathbf{W}}_u$ associated with the tangent space $T_u \mathbf{V}^{2n}$, for $u \in \mathbf{V}^{2n}$. It should be noted that the fibre product (33) can be trivially extended to the space of $\tilde{\mathbf{W}}$ -valued N-adapted differential forms $\tilde{\mathcal{W}} \otimes \Lambda$ by means of the usual exterior product of the scalar forms Λ , where $\tilde{\mathcal{W}}$ denotes the sheaf of smooth sections of $\tilde{\mathbf{W}}$. There is a standard grading on Λ denoted $\deg_a$. One also introduces gradings $\deg_v, \deg_s, \deg_a$ on $\tilde{\mathcal{W}} \otimes \Lambda$ defined on homogeneous elements $v, z^\alpha, \check{e}^\alpha$ as follows: $\deg_v(v) = 1$, $\deg_s(z^\alpha) = 1$, $\deg_a(\check{e}^\alpha) = 1$, and all other gradings of the elements $v, z^\alpha, \check{e}^\alpha$ are set to zero. In this case, the product $\circ$ from (33) on $\tilde{\mathcal{W}} \otimes \Lambda$ is bigraded. This is written w.r.t the grading $Deg = 2\deg_v + \deg_s$ and the grading $\deg_a$.

4.1 Normal Fedosov's d-operators

The normal d-connection $\hat{\mathbf{D}} = \{\hat{\Gamma}_{\alpha\beta}^\gamma\}$ (24) can be extended to operators

$$\hat{\mathbf{D}}(a \otimes \lambda) \doteq \left(\check{e}_\alpha(a) - u^\beta \hat{\Gamma}_{\alpha\beta}^\gamma z^\gamma \check{e}_\alpha(a) \right) \otimes (\check{e}^\alpha \wedge \lambda) + a \otimes d\lambda, \quad (35)$$

on $\tilde{\mathcal{W}} \otimes \Lambda$, where $z^\gamma \check{e}_\alpha$ is $\check{e}_\alpha$ redefined in z -variables. This operator $\hat{\mathbf{D}}$ is a N-adapted $\deg_a$ -graded derivation of the distinguished algebra $(\tilde{\mathcal{W}} \otimes \Lambda, \circ)$, called d-algebra. Such a property follows from (33) and (35)).

Definition 4.1 *The Fedosov distinguished operators (d-operators) $\check{\delta}$ and $\check{\delta}^{-1}$ on $\tilde{\mathcal{W}} \otimes \Lambda$, are defined*

$$\check{\delta}(a) = \check{e}^\alpha \wedge z^\alpha \check{e}_\alpha(a), \text{ and } \check{\delta}^{-1}(a) = \begin{cases} \frac{i}{p+q} z^\alpha \check{e}_\alpha(a), & \text{if } p+q > 0, \\ 0, & \text{if } p+q = 0, \end{cases} \quad (36)$$

where any $a \in \tilde{\mathcal{W}} \otimes \Lambda$ is homogeneous w.r.t. the grading $\deg_s$ and $\deg_a$ with $\deg_s(a) = p$ and $\deg_a(a) = q$.

The d-operators (36) define the formula $a = (\check{\delta} \check{\delta}^{-1} + \check{\delta}^{-1} \check{\delta} + \sigma)(a)$, where $a \mapsto \sigma(a)$ is the projection on the $(\deg_s, \deg_a)$ -bihomogeneous part of a of degree zero, $\deg_s(a) = \deg_a(a) = 0$; $\check{\delta}$ is also a $\deg_a$ -graded derivation of the d-algebra $(\check{\mathcal{W}} \otimes \mathbf{\Lambda}, \circ)$. In order to emphasize the almost Kähler structure, we used the canonical almost symplectic geometric objects defined by a fixed $\mathcal{L}$. Nevertheless, we can always change the "Lagrangian mechanics variables" and redefine $\check{\theta}$, $\check{\mathbf{e}}_\alpha$ and $\hat{\mathbf{T}}_{\alpha\beta}^\gamma$ with respect to arbitrary frame and co-frame bases using vielbeins $\mathbf{e}_\alpha^\underline{a}$ and their duals $\mathbf{e}_\alpha^\alpha$, defined by $e^i_{\underline{i}}, e^a_{\underline{i}}$ and $e^a_{\underline{a}}$ satisfying (7).

We can provide a "N-adapted" proof [33, 2] of

Proposition 4.1 *The torsion and curvature canonical d-operators of the extension of $\hat{\mathbf{D}}$ to $\check{\mathcal{W}} \otimes \mathbf{\Lambda}$, are computed*

$${}^z\hat{\mathcal{T}} \doteq \frac{z^\gamma}{2} \check{\theta}_{\gamma\tau} \hat{\mathbf{T}}_{\alpha\beta}^\tau(u) \check{\mathbf{e}}^\alpha \wedge \check{\mathbf{e}}^\beta, \quad (37)$$

and

$${}^z\hat{\mathcal{R}} \doteq \frac{z^\gamma z^\varphi}{4} \check{\theta}_{\gamma\tau} \hat{\mathbf{R}}_{\varphi\alpha\beta}^\tau(u) \check{\mathbf{e}}^\alpha \wedge \check{\mathbf{e}}^\beta, \quad (38)$$

where the nontrivial coefficients of $\hat{\mathbf{T}}_{\alpha\beta}^\tau$ and $\hat{\mathbf{R}}_{\varphi\alpha\beta}^\tau$ are defined respectively by formulas (29) and (31).

By straightforward verifications, it follows the proof of

Theorem 4.1 *The properties*

$$[\hat{\mathbf{D}}, \check{\delta}] = \frac{i}{v} ad_{Wick}({}^z\hat{\mathcal{T}}) \text{ and } \hat{\mathbf{D}}^2 = -\frac{i}{v} ad_{Wick}({}^z\hat{\mathcal{R}}), \quad (39)$$

hold for the above operators, where $[\cdot, \cdot]$ is the $\deg_a$ -graded commutator of endomorphisms of $\check{\mathcal{W}} \otimes \mathbf{\Lambda}$ and ad_{Wick} is defined via the $\deg_a$ -graded commutator in $(\check{\mathcal{W}} \otimes \mathbf{\Lambda}, \circ)$.

The formulas (39) can be redefined for any linear connection structure on $\mathbf{V}^{2n}$. For example, we consider how similar formulas can be provided for the Levi-Civita connection.

4.2 Fedosov's d-operators and the Levi-Civita connection

For any metric structure $\mathbf{g}$ on a manifold $\mathbf{V}^{2n}$, the Levi-Civita connection $\nabla = \{ \cdot \Gamma_{\beta\gamma}^\alpha \}$ is by definition the unique linear connection that is metric compatible ($\nabla g = 0$) and torsionless ($\cdot \mathcal{T} = 0$). It is not a d-connection because

it does not preserve the N-connection splitting under parallel transports (10). Let us parametrize its coefficients in the form

$$\begin{aligned} \Gamma_{\beta\gamma}^\alpha &= ({}_1L_{jk}^i, {}_1L_{jk}^a, {}_1L_{bk}^i, {}_1L_{bk}^a, {}_1C_{jb}^i, {}_1C_{jb}^a, {}_1C_{bc}^i, {}_1C_{bc}^a), \text{ where} \\ \nabla_{\check{e}_k}(\check{e}_j) &= {}_1L_{jk}^i\check{e}_i + {}_1L_{jk}^ae_a, \quad \nabla_{\check{e}_k}(e_b) = {}_1L_{bk}^i\check{e}_i + {}_1L_{bk}^ae_a, \\ \nabla_{e_b}(\check{e}_j) &= {}_1C_{jb}^i\check{e}_i + {}_1C_{jb}^ae_a, \quad \nabla_{e_c}(e_b) = {}_1C_{bc}^i\check{e}_i + {}_1C_{bc}^ae_a. \end{aligned}$$

A straightforward calculation shows that the coefficients of the Levi-Civita connection can be expressed as

$$\begin{aligned} {}_1L_{jk}^a &= -\widehat{C}_{jb}^i\check{g}_{ik}\check{g}^{ab} - \frac{1}{2}\check{\Omega}_{jk}^a, \quad {}_1L_{bk}^i = \frac{1}{2}\check{\Omega}_{jk}^c\check{g}_{cb}\check{g}^{ji} - \Xi_{jk}^{ih}\widehat{C}_{hb}^j, \\ {}_1L_{jk}^i &= \widehat{L}_{jk}^i, \quad {}_1L_{bk}^a = \widehat{L}_{bk}^a + {}^\pm\Xi_{cd}^{ab} \circ L_{bk}^c, \quad {}_1C_{kb}^i = \widehat{C}_{kb}^i + \frac{1}{2}\check{\Omega}_{jk}^a\check{g}_{cb}\check{g}^{ji} + \Xi_{jk}^{ih}\widehat{C}_{hb}^j, \\ {}_1C_{jb}^a &= -{}^\pm\Xi_{cb}^{ad} \circ L_{dj}^c, \quad {}_1C_{bc}^a = \widehat{C}_{bc}^a, \quad {}_1C_{ab}^i = -\frac{\check{g}^{ij}}{2} \{ \circ L_{aj}^c\check{g}_{cb} + \circ L_{bj}^c\check{g}_{ca} \}, \end{aligned} \quad (40)$$

where $e_b = \partial/\partial y^a$, $\check{\Omega}_{jk}^a$ are computed as in (15) but for the canonical N-connection $\check{\mathbf{N}}$ (5), $\Xi_{jk}^{ih} = \frac{1}{2}(\delta_j^i\delta_k^h - \check{g}_{jk}\check{g}^{ih})$, ${}^\pm\Xi_{cd}^{ab} = \frac{1}{2}(\delta_c^a\delta_d^b \pm \check{g}_{cd}\check{g}^{ab})$, $\circ L_{aj}^c = \widehat{L}_{aj}^c - e_a(\check{N}_j^c)$, $\check{g}_{ik}$ and $\check{g}^{ab}$ are defined for the representation of the metric in Lagrange-Finsler variables (3) and the normal d-connection $\widehat{\Gamma}_{\beta\gamma}^\alpha = (\widehat{L}_{jk}^i, {}^v\widehat{C}_{bc}^a)$ (24) is given by coefficients (25).

Let introduce the distortion d-tensor ${}_1Z_{\alpha\beta}^\gamma$ with N-adapted coefficients

$$\begin{aligned} {}_1Z_{jk}^a &= -\widehat{C}_{jb}^i\check{g}_{ik}\check{g}^{ab} - \frac{1}{2}\check{\Omega}_{jk}^a, \quad {}_1Z_{bk}^i = \frac{1}{2}\check{\Omega}_{jk}^c\check{g}_{cb}\check{g}^{ji} - \Xi_{jk}^{ih}\widehat{C}_{hb}^j, \\ {}_1Z_{jk}^i &= 0, \quad {}_1Z_{bk}^a = {}^\pm\Xi_{cd}^{ab} \circ L_{bk}^c, \quad {}_1Z_{kb}^i = \frac{1}{2}\check{\Omega}_{jk}^a\check{g}_{cb}\check{g}^{ji} + \Xi_{jk}^{ih}\widehat{C}_{hb}^j, \\ {}_1Z_{jb}^a &= -{}^\pm\Xi_{cb}^{ad} \circ L_{dj}^c, \quad {}_1Z_{bc}^a = 0, \quad {}_1Z_{ab}^i = -\frac{\check{g}^{ij}}{2} [\circ L_{aj}^c\check{g}_{cb} + \circ L_{bj}^c\check{g}_{ca}], \end{aligned} \quad (41)$$

The next result follows from the above arguments.

Proposition 4.2 *The N-adapted coefficients, of the normal d-connection and of the distortion d-tensors define the Levi-Civita connection as*

$${}_1\Gamma_{\alpha\beta}^\gamma = \widehat{\Gamma}_{\alpha\beta}^\gamma + {}_1Z_{\alpha\beta}^\gamma, \quad (42)$$

where ${}_1Z_{\alpha\beta}^\gamma$ are given by formulas (42) and h- and v-components of $\widehat{\Gamma}_{\beta\gamma}^\alpha$ are given by (25).

We emphasize that all components of ${}_1\Gamma_{\alpha\beta}^\gamma, \widehat{\Gamma}_{\alpha\beta}^\gamma$ and ${}_1Z_{\alpha\beta}^\gamma$ are uniquely defined by the coefficients of d-metric (1), or (equivalently) by (3) and (5). The constructions can be obtained for any $n + n$ splitting on V^{2n} , which for suitable $\mathcal{L}$, or $\mathcal{F}$, admit a Lagrange, or Finsler, like representation of geometric objects.

By proposition 4.1, the expressions for the curvature and torsion of canonical d-operators of the extension of ∇ to $\check{\mathcal{W}} \otimes \Lambda$, are

$$\begin{aligned} {}^z\mathcal{R} &\doteq \frac{{}^z\gamma {}^z\varphi}{4} \check{\theta}_{\gamma\tau} {}_1R_{\varphi\alpha\beta}^\tau(u) \check{\mathbf{e}}^\alpha \wedge \check{\mathbf{e}}^\beta, \\ {}^z\mathcal{T} &\doteq \frac{{}^z\gamma}{2} \check{\theta}_{\gamma\tau} {}_1T_{\alpha\beta}^\tau(u) \check{\mathbf{e}}^\alpha \wedge \check{\mathbf{e}}^\beta \equiv 0, \end{aligned} \quad (43)$$

where ${}_1T_{\alpha\beta}^\tau = 0$, by definition, and ${}_1R_{\varphi\alpha\beta}^\tau$ is computed with respect to the N-adapted Lagange–Finsler canonical bases by introducing $\widehat{\Gamma}_{\alpha\beta}^\gamma = -{}_1\Gamma_{\alpha\beta}^\gamma + {}_1Z_{\alpha\beta}^\gamma$, see (42), into (31). To the N-adapted d-operator (35), we can associate

$$\widehat{\nabla}(a \otimes \lambda) \doteq \left(\check{\mathbf{e}}_\alpha(a) - u^\beta {}_1\Gamma_{\alpha\beta}^\gamma {}^z\check{\mathbf{e}}_\alpha(a) \right) \otimes (\check{\mathbf{e}}^\alpha \wedge \lambda) + a \otimes d\lambda, \quad (44)$$

on $\check{\mathcal{W}} \otimes \Lambda$, where ${}^z\check{\mathbf{e}}_\alpha$ is $\check{\mathbf{e}}_\alpha$ redefined in z -variables. This almost symplectic connection $\widehat{\nabla}$ is torsionless and, in general, is not adapted to the N-connection structures.

Corollary 4.1 *For the Levi–Civita connection $\nabla = \{ {}_1\Gamma_{\beta\gamma}^\alpha \}$ on a N-anholonomic manifold $\mathbf{V}^{2n}$, we have:*

$$\left[\widehat{\nabla}, \check{\delta} \right] = 0 \text{ and } \widehat{\nabla}^2 = -\frac{i}{v} ad_{Wick}({}^z\mathcal{R}),$$

where $\widehat{\nabla}$ is defined by formula (44), ${}^z\mathcal{R}$ is given by (43), $[\cdot, \cdot]$ is the $\deg_a$ -graded commutator of endomorphisms of $\check{\mathcal{W}} \otimes \Lambda$ and ad_{Wick} is defined via the $\deg_a$ -graded commutator in $(\check{\mathcal{W}} \otimes \Lambda, \circ)$.

Proof. It is a straightforward consequence of the Theorem 4.1 for the Levi–Civita and curvature operators extended on $\check{\mathcal{W}} \otimes \Lambda$. $\square$

Prescribing a $n + n$ splitting on $\mathbf{V}^{2n}$, we can work equivalently with any metric compatible linear connection structure which is N-adapted, or not, if such a connection is completely defined by the (pseudo) Riemannian metric structure. It is preferable to use the approach with the normal d-connection because this way we have both an almost symplectic analogy and Lagrange,

or Finsler, like interpretation of geometric objects. In standard classical gravity, in order to solve some physical problems, it is more convenient to work with the Levi–Civita connection or its spin like representations (for instance, in the Einstein–Dirac theory). The self–dual and further generalizations to Ashtekar variables are more convenient, respectively, in canonical ADN classical and quantum gravity and/or loop quantum gravity.

It should be noted that the formulas for Fedosov’s d –operators and their properties do not depend in explicit form on generating functions $\mathcal{L}$, or $\mathcal{F}$. Such a function may be formally introduced for elaborating a Lagrange mechanics, or Finsler, modeling for a (pseudo) Riemannian space with a general $n + n$ nonholonomic splitting. This way, we emphasize that the Fedosov’s approach is valid for various type of (pseudo) Riemann, Riemann–Cartan, Lagrange–Finsler, almost Kähler and other types of holonomic and non-holonomic manifolds used for geometrization of mechanical and field models. Nevertheless, the constructions are performed in a general form and the final results do not depend on any ”background” structures. We conclude that $3 + 1$ fibration approaches are more natural for loop quantum gravity, but the models with nonholonomic $2 + 2$ splitting result in almost Kähler quantum models; although both types of quantization, loop and deformation, provide background independent constructions.

5 Deformation Quantization of Einstein and Lagrange Spaces

Formulating a (pseudo) Riemannian geometry in Lagrange–Finsler variables, we can quantize the metric, frame and linear connection structures following standard methods for deformation quantization of almost Kähler manifolds. The goal of this section is to provide the main Fedosov type results for such constructions and to show how the Einstein manifolds can be encoded into the topological structure of such quantized nonholonomic spaces.

5.1 Fedosov’s theorems for normal d –connections

The third main result of this work will be stated below by three theorems for the normal d –connection (equivalently, canonical almost symplectic structure) $\widehat{\mathbf{D}} \equiv {}_{\theta}\widehat{\mathbf{D}}$ (24). Such results were formulated originally in Fedosov’s works [5, 6] and generalized, for instance, for various classes of metric compatible affine connections, canonical Lagrange–Finsler connections and effective locally anisotropic quantum gravities, see Refs. [33, 1, 2, 3].

Theorem 5.1 Any (pseudo) Riemannian metric $\mathbf{g}$ (1) (equivalently, $\mathbf{g} = \check{\mathbf{g}}$ (3)) defines a flat normal Fedosov d -connection

$$\widehat{\mathcal{D}} \doteq -\check{\delta} + \widehat{\mathbf{D}} - \frac{i}{v} \text{ad}_{\text{Wick}}(r)$$

satisfying the condition $\widehat{\mathcal{D}}^2 = 0$, where the unique element $r \in \check{\mathcal{W}} \otimes \mathbf{\Lambda}$, $\deg_a(r) = 1$, $\check{\delta}^{-1}r = 0$, solves the equation

$$\check{\delta}r = \widehat{\mathcal{T}} + \widehat{\mathcal{R}} + \widehat{\mathbf{D}}r - \frac{i}{v} r \circ r$$

and this element can be computed recursively with respect to the total degree Deg as follows:

$$\begin{aligned} r^{(0)} &= r^{(1)} = 0, r^{(2)} = \check{\delta}^{-1}\widehat{\mathcal{T}}, r^{(3)} = \check{\delta}^{-1}\left(\widehat{\mathcal{R}} + \widehat{\mathbf{D}}r^{(2)} - \frac{i}{v}r^{(2)} \circ r^{(2)}\right), \\ r^{(k+3)} &= \check{\delta}^{-1}\left(\widehat{\mathbf{D}}r^{(k+2)} - \frac{i}{v}\sum_{l=0}^k r^{(l+2)} \circ r^{(l+2)}\right), k \geq 1, \end{aligned}$$

where by $a^{(k)}$ we denoted the Deg -homogeneous component of degree k of an element $a \in \check{\mathcal{W}} \otimes \mathbf{\Lambda}$.

Proof. It follows from straightforward verifications of the property $\widehat{\mathcal{D}}^2 = 0$ using for r formal series of type (34) and the formulas for N -adapted coefficients: (25) for $\widehat{\mathbf{D}}$, (29) for $\widehat{\mathcal{T}}$, (31) for $\widehat{\mathcal{R}}$, and the properties of Fedosov's d -operators (36) stated by Theorem 4.1. The length of this paper does not allow us to present such a tedious calculation which is a N -adapted version for corresponding "hat" operators, see the related work in [5, 6, 33]. $\square$

The procedure of deformation quantization is related to the definition of a star-product which in our approach can be defined canonically as in [33] because the normal d -connection $\widehat{\mathbf{D}}$ is a N -adapted variant of the affine and almost symplectic connection considered in that work. This provides a proof for

Theorem 5.2 A star-product on the almost Kähler model of a (pseudo) Riemannian space in Lagrange-Finsler variables is defined on $C^\infty(\mathbf{V}^{2n})[[v]]$ by formula

$${}^1f * {}^2f \doteq \sigma(\tau({}^1f)) \circ \sigma(\tau({}^2f)),$$

where the projection $\sigma : \check{\mathcal{W}}_{\widehat{\mathcal{D}}} \rightarrow C^\infty(\mathbf{V}^{2n})[[v]]$ onto the part of $\deg_s$ -degree zero is a bijection and the inverse map $\tau : C^\infty(\mathbf{V}^{2n})[[v]] \rightarrow \check{\mathcal{W}}_{\widehat{\mathcal{D}}}$ can be calculated recursively w.r.t the total degree Deg ,

$$\begin{aligned}\tau(f)^{(0)} &= f \text{ and, for } k \geq 0, \\ \tau(f)^{(k+1)} &= \check{\delta}^{-1} \left(\widehat{\mathbf{D}}\tau(f)^{(k)} - \frac{i}{v} \sum_{l=0}^k \text{ad}_{\text{Wick}}(r^{(l+2)})(\tau(f)^{(k-l)}) \right).\end{aligned}$$

We denote by ${}^f\xi$ the Hamiltonian vector field corresponding to a function $f \in C^\infty(\mathbf{V}^{2n})$ on space $(\mathbf{V}^{2n}, \check{\theta})$ and consider the antisymmetric part ${}^{-1}C({}^1f, {}^2f) \doteq \frac{1}{2} (C({}^1f, {}^2f) - C({}^2f, {}^1f))$ of bilinear operator $C({}^1f, {}^2f)$. We say that a star-product (32) is normalized if ${}^{-1}C({}^1f, {}^2f) = \frac{i}{2} \{ {}^1f, {}^2f \}$, where $\{\cdot, \cdot\}$ is the Poisson bracket. For the normalized $*$, the bilinear operator ${}^{-2}C$ defines a de Rham–Chevalley 2-cocycle, when there is a unique closed 2-form $\check{\varkappa}$ such that

$${}^{-2}C({}^1f, {}^2f) = \frac{1}{2} \check{\varkappa}({}^{f_1}\xi, {}^{f_2}\xi) \quad (45)$$

for all ${}^1f, {}^2f \in C^\infty(\mathbf{V}^{2n})$. This is used to introduce $c_0(*) \doteq [\check{\varkappa}]$ as the equivalence class.

A straightforward computation of ${}^{-2}C$ from (45) and the results of Theorem 5.2 provide the proof of

Lemma 5.1 *The unique 2-form defined by the normal d-connection can be computed as*

$$\check{\varkappa} = -\frac{i}{8} \check{\mathbf{J}}_{\tau}{}^{\alpha'} \widehat{\mathcal{R}}^{\tau}{}_{\alpha'} - \frac{i}{6} d \left(\check{\mathbf{J}}_{\tau}{}^{\alpha'} \widehat{\mathbf{T}}^{\tau}{}_{\alpha'\beta} \check{\mathbf{e}}^{\beta} \right),$$

where the coefficients of the curvature and torsion 2-forms of the normal d-connection 1-form are given respectively by formulas (30) and (28).

We now define another canonical class $\check{\varepsilon}$, for ${}^{\check{N}}T\mathbf{V}^{2n} = h\mathbf{V}^{2n} \oplus v\mathbf{V}^{2n}$, where the left label indicates that the tangent bundle is split nonholonomically by the canonical N-connection structure $\check{\mathbf{N}}$. We can perform a distinguished complexification of such second order tangent bundles in the form $T_{\mathbb{C}}({}^{\check{N}}T\mathbf{V}^{2n}) = T_{\mathbb{C}}(h\mathbf{V}^{2n}) \oplus T_{\mathbb{C}}(v\mathbf{V}^{2n})$ and introduce $\check{\varepsilon}$ as the first Chern class of the distributions $T'_{\mathbb{C}}({}^N T\mathbf{V}^{2n}) = T'_{\mathbb{C}}(h\mathbf{V}^{2n}) \oplus T'_{\mathbb{C}}(v\mathbf{V}^{2n})$ of couples of vectors of type $(1, 0)$ both for the h- and v-parts. In explicit form, we can calculate $\check{\varepsilon}$ by using the d-connection $\widehat{\mathbf{D}}$ and the h- and v-projections $h\Pi = \frac{1}{2}(Id_h - iJ_h)$ and $v\Pi = \frac{1}{2}(Id_v - iJ_v)$, where Id_h and Id_v are respective

identity operators and J_h and J_v are almost complex operators, which are projection operators onto corresponding $(1, 0)$ -subspaces. Introducing the matrix $(h\Pi, v\Pi) \widehat{\mathcal{R}} (h\Pi, v\Pi)^T$, where $(\dots)^T$ means transposition, as the curvature matrix of the N-adapted restriction of of the normal d-connection $\widehat{\mathbf{D}}$ to $T'_\mathbb{C} \left(\tilde{N}T\mathbf{V}^{2n} \right)$, we compute the closed Chern–Weyl form

$$\tilde{\gamma} = -iTr \left[(h\Pi, v\Pi) \widehat{\mathcal{R}} (h\Pi, v\Pi)^T \right] = -iTr \left[(h\Pi, v\Pi) \widehat{\mathcal{R}} \right] = -\frac{1}{4} \check{\mathbf{J}}_\tau^{\alpha'} \widehat{\mathcal{R}}_\tau^{\alpha'}. \quad (46)$$

We get that the canonical class is $\tilde{\varepsilon} \doteq [\tilde{\gamma}]$, which proves the

Theorem 5.3 *The zero-degree cohomology coefficient $c_0(*)$ for the almost Kähler model of a (pseudo) Riemannian space defined by d-tensor $\mathbf{g}$ (1) (equivalently, by $\check{\mathbf{g}}$ (3)) is computed $c_0(*) = -(1/2i) \tilde{\varepsilon}$.*

The coefficient $c_0(*)$ can be similarly computed for the case when a metric of type (1) is a solution of the Einstein equations and this zero-degree coefficient defines certain quantum properties of the gravitational field. A more rich geometric structure should be considered if we define a value similar to $c_0(*)$ encoding the information about Einstein manifolds deformed into corresponding quantum configurations.

5.2 The zero-degree cohomology coefficient for Einstein manifolds

The priority of deformation quantization is that we can elaborate quantization schemes when metric, vielbein and connection fields are not obligatory subjected to satisfy certain field equations and/or derived by a variational procedure. For instance, such geometric and/or BRST quantization approaches were proposed in Ref. [36, 37]. On the other hand, in certain canonical and loop quantization models, the gravitational field equations are considered as the starting point for deriving a quantization formalism. In such cases, the Einstein equations are expressed into "lapse" and "shift" (and/or generalized Ashtekar) variables and the quantum variant of the gravitational field equations is prescribed to be in the form of Wheeler De Witt equations (or corresponding systems of constraints in complex/real generalized connection and dreibein variables). In this section, we analyze the problem of encoding the Einstein equations into a geometric formalism of nonholonomic deformation quantization.

5.2.1 Gravitational field equations

For any d-connection $\mathbf{D} = \{\mathbf{\Gamma}\}$, we can define the Ricci tensor $Ric(\mathbf{D}) = \{\mathbf{R}_{\beta\gamma} \doteq \mathbf{R}_{\beta\gamma\alpha}^{\alpha}\}$ and the scalar curvature ${}^sR \doteq \mathbf{g}^{\alpha\beta}\mathbf{R}_{\alpha\beta}$ ($\mathbf{g}^{\alpha\beta}$ being the inverse matrix to $\mathbf{g}_{\alpha\beta}$ (1)). If a d-connection is uniquely determined by a metric in a unique metric compatible form, $\mathbf{D}\mathbf{g} = 0$, (in general, the torsion of $\mathbf{D}$ is not zero, but induced canonically by the coefficients of $\mathbf{g}$), we can postulate in straightforward form the field equations

$$\mathbf{R}_{\beta}^{\alpha} - \frac{1}{2}({}^sR + \lambda)\mathbf{e}_{\beta}^{\alpha} = 8\pi G\mathbf{T}_{\beta}^{\alpha}, \quad (47)$$

where $\mathbf{T}_{\beta}^{\alpha}$ is the effective energy-momentum tensor, λ is the cosmological constant, G is the Newton constant in the units when the light velocity $c = 1$, and $\mathbf{e}_{\beta}^{\alpha} = \mathbf{e}_{\beta}^{\alpha}\partial/\partial u^{\alpha}$ is the N-elongated operator (11).

Let us consider the absolute antisymmetric tensor $\epsilon_{\alpha\beta\gamma\delta}$ and effective source 3-form

$$\mathcal{T}_{\beta} = \mathbf{T}_{\beta}^{\alpha} \epsilon_{\alpha\beta\gamma\delta} du^{\beta} \wedge du^{\gamma} \wedge du^{\delta}$$

and express the curvature tensor $\mathcal{R}_{\gamma}^{\tau} = \mathbf{R}_{\gamma\alpha\beta}^{\tau} \mathbf{e}^{\alpha} \wedge \mathbf{e}^{\beta}$ of $\mathbf{\Gamma}_{\beta\gamma}^{\alpha} = {}_1\mathbf{\Gamma}_{\beta\gamma}^{\alpha} - Z_{\beta\gamma}^{\alpha}$ as $\mathcal{R}_{\gamma}^{\tau} = {}_1\mathcal{R}_{\gamma}^{\tau} - \mathcal{Z}_{\gamma}^{\tau}$, where ${}_1\mathcal{R}_{\gamma}^{\tau} = {}_1R_{\gamma\alpha\beta}^{\tau} \mathbf{e}^{\alpha} \wedge \mathbf{e}^{\beta}$ is the curvature 2-form of the Levi-Civita connection ∇ and the distortion of curvature 2-form $\mathcal{Z}_{\gamma}^{\tau}$ is defined by $Z_{\beta\gamma}^{\alpha}$. For the gravitational $(\mathbf{e}, \mathbf{\Gamma})$ and matter ϕ fields, we consider the effective action

$$S[\mathbf{e}, \mathbf{\Gamma}, \phi] = {}^{gr}S[\mathbf{e}, \mathbf{\Gamma}] + {}^{matter}S[\mathbf{e}, \mathbf{\Gamma}, \phi].$$

Theorem 5.4 *The equations (47) can be represented as 3-form equations*

$$\epsilon_{\alpha\beta\gamma\tau} \left(\mathbf{e}^{\alpha} \wedge \mathcal{R}^{\beta\gamma} + \lambda \mathbf{e}^{\alpha} \wedge \mathbf{e}^{\beta} \wedge \mathbf{e}^{\gamma} \right) = 8\pi G \mathcal{T}_{\tau} \quad (48)$$

following from the action by varying the components of $\mathbf{e}_{\beta}$, when

$$\begin{aligned} \mathcal{T}_{\tau} &= {}^m\mathcal{T}_{\tau} + {}^Z\mathcal{T}_{\tau}, \\ {}^m\mathcal{T}_{\tau} &= {}^m\mathbf{T}_{\tau}^{\alpha} \epsilon_{\alpha\beta\gamma\delta} du^{\beta} \wedge du^{\gamma} \wedge du^{\delta}, \\ {}^Z\mathcal{T}_{\tau} &= (8\pi G)^{-1} Z_{\tau}^{\alpha} \epsilon_{\alpha\beta\gamma\delta} du^{\beta} \wedge du^{\gamma} \wedge du^{\delta}, \end{aligned}$$

where ${}^m\mathbf{T}_{\tau}^{\alpha} = \delta {}^{matter}S / \delta \mathbf{e}_{\tau}^{\alpha}$ are equivalent to the usual Einstein equations for the Levi-Civita connection ∇ ,

$${}_1\mathbf{R}_{\beta}^{\alpha} - \frac{1}{2}({}_1{}^sR + \lambda)\mathbf{e}_{\beta}^{\alpha} = 8\pi G {}^m\mathbf{T}_{\beta}^{\alpha}.$$

Proof. It is a usual textbook and/or differential form calculus (see, for instance, [16, 9]), but with respect to N -adapted bases (11) and (12) for a metric compatible d -connection (47). $\square$

As a particular case, for the Einstein gravity in Lagrange–Finsler variables, we obtain:

Corollary 5.1 *The vacuum Einstein equations, with cosmological constant in terms of the canonical N -adapted vierbeins and normal d -connection, are*

$$\epsilon_{\alpha\beta\gamma\tau} \left(\check{\mathbf{e}}^\alpha \wedge \widehat{\mathcal{R}}^{\beta\gamma} + \lambda \check{\mathbf{e}}^\alpha \wedge \check{\mathbf{e}}^\beta \wedge \check{\mathbf{e}}^\gamma \right) = 8\pi G \, {}^Z\widehat{\mathcal{T}}_\tau, \quad (49)$$

or, in terms of the Levi–Civita connection

$$\epsilon_{\alpha\beta\gamma\tau} \left(\check{\mathbf{e}}^\alpha \wedge {}_i\mathcal{R}^{\beta\gamma} + \lambda \check{\mathbf{e}}^\alpha \wedge \check{\mathbf{e}}^\beta \wedge \check{\mathbf{e}}^\gamma \right) = 0.$$

Proof. The conditions of the mentioned Theorem 5.4 are redefined for the co-frames $\check{\mathbf{e}}^\alpha$ elongated by the canonical N -connection (5), deformation of linear connections (42) and curvature (31) with deformation of curvature 2-form of type

$$\widehat{\mathcal{R}}^\tau_\gamma = {}_i\mathcal{R}^\tau_\gamma - \widehat{\mathcal{Z}}^\tau_\gamma. \quad (50)$$

We put "hat" on ${}^Z\widehat{\mathcal{T}}_\tau$ because this value is computed using the normal d -connection. $\square$

Using formulas (49) and (50), we can write

$$\widehat{\mathcal{R}}^{\beta\gamma} = -\lambda \check{\mathbf{e}}^\beta \wedge \check{\mathbf{e}}^\gamma - \widehat{\mathcal{Z}}^{\beta\gamma} \text{ and } {}_i\mathcal{R}^{\beta\gamma} = -\lambda \check{\mathbf{e}}^\beta \wedge \check{\mathbf{e}}^\gamma. \quad (51)$$

Such formulas are necessary for encoding the vacuum field equations into the cohomological structure of quantum almost Kähler model of Einstein gravity.

5.2.2 The Chern–Weyl form and Einstein equations

Introducing the formulas (49) and (51) into the conditions of Lemma 5.1 and Theorem 5.3, we obtain the forth main result in this work:

Theorem 5.5 *The zero-degree cohomology coefficient $c_0(*)$ for the almost Kähler model of an Einstein space defined by a d -tensor $\mathbf{g}$ (1) (equivalently, by $\check{\mathbf{g}}$ (3)) as a solution of (49) is $c_0(*) = -(1/2i) \check{\mathcal{E}}$, for $\check{\mathcal{E}} \doteq [\check{\gamma}]$, where*

$$\check{\gamma} = \frac{1}{4} \check{\mathbf{J}}_{\tau\alpha} \left(-\lambda \check{\mathbf{e}}^\tau \wedge \check{\mathbf{e}}^\alpha + \widehat{\mathcal{Z}}^{\tau\alpha} \right). \quad (52)$$

Proof. We sketch the key points of the proof which follows from (46) and (51). It should be noted that for $\lambda = 0$ the 2-form $\widehat{\mathcal{Z}}^{\tau\alpha}$ is defined by the deformation d-tensor from the Levi-Civita connection to the normal d-connection (42), see formulas (41). Such objects are defined by classical vacuum solutions of the Einstein equations. We conclude that $c_0(*)$ encodes the vacuum Einstein configurations, in general, with nontrivial constants and their quantum deformations. $\square$

If the Wheeler De Witt equations represent a quantum version of the Einstein equations for loop quantum gravity (see discussions in Refs. [9, 10, 11]), the Chern–Weyl 2-form (52) can be used to define the quantum version of Einstein equations (48) in the deformation quantization approach:

Corollary 5.2 *In Lagrange–Finsler variables, the quantum field equations corresponding to Einstein’s general relativity are*

$$\check{\mathbf{e}}^\alpha \wedge \check{\gamma} = \epsilon^{\alpha\beta\gamma\tau} 2\pi G \check{\mathbf{J}}_{\beta\gamma} \widehat{\mathcal{T}}_\tau - \frac{\lambda}{4} \check{\mathbf{J}}_{\beta\gamma} \check{\mathbf{e}}^\alpha \wedge \check{\mathbf{e}}^\beta \wedge \check{\mathbf{e}}^\gamma. \quad (53)$$

Proof. Multiplying $\check{\mathbf{e}}^\alpha \wedge$ to (52) written in Lagrange–Finsler variables and taking into account (48), re-written also in the form adapted to the canonical N-connection, and introducing the almost complex operator $\check{\mathbf{J}}_{\beta\gamma}$, we get the almost symplectic form of Einstein’s equations (53). $\square$

It should be noted that even in the vacuum case, when $\lambda = 0$, the 2-form $\check{\gamma}$ (52) from (53) is not zero but defined by $\widehat{\mathcal{T}}_\tau = {}^Z\widehat{\mathcal{T}}_\tau$.

Finally, we emphasize that an explicit computation of $\check{\gamma}$ for nontrivial matter fields has yet to be performed for a deformation quantization model in which interacting gravitational and matter fields are geometrized in terms of an almost Kähler model defined for spinor and fiber bundles on spacetime. This is a subject for further investigations.

6 Conclusions and Discussion

So far we have dealt with the deformation quantization of general relativity in Lagrange–Finsler variables, inducing a nonholonomic 2+2 splitting, as a geometric alternative to the 3+1 setting to loop quantum gravity. In our approach, the methods of geometric mechanics and Finsler geometry are canonically combined in order to convert general relativity into an almost Kähler structure for which a well defined formalism of geometric quantization exists. The formalism is elaborated in a background independent and

nonperturbative form, for Lorentzian gravitational fields on four dimensional manifolds, with possible extensions to extra dimensions.

There are many physical interesting formulations of gravity theories in differential form, with tetrad, spinor and different connection variables. For instance, the 3+1 splitting and Ashtekar variables resulted in a similarity with Yang Mills theory and allowed us to simplify the constraints and provide consistent loop quantum gravity formulations. In order to construct almost Kähler models of classical and quantum gravity, it is more convenient to use a 2+2 splitting with prescribed nonholonomic frame structures when the metric and linear connection transform into canonical symplectic forms and connections. The main advantage of this approach is that we do not have to solve constraint equations and can apply directly the methods of deformation quantization. We can work both with tetrad and connection variables, and the methods can be generalized for Lagrange and Finsler spaces, almost Kähler and almost Poisson structures, nonsymmetric metrics and noncommutative geometries.

The facts that Einstein's theory is diffeomorphic invariant and preserves local Lorentz invariance are crucial features at the classical level and provide strong motivations to preserve such symmetries at the quantum level. We emphasize here that by prescribing a distribution by defining a non-holonomic frame structure with associated nonlinear connection on a (semi) Riemannian manifold, we do not break general covariance. We chose to work with a class of frame transforms that did not affect the general properties of classical and quantum gravity theories. In general, all constructions can be re-defined for arbitrary frames and coordinate systems.

Let us outline the main results (four) of this paper. The first one is provided by Theorem 2.1, stating that any (pseudo) Riemannian space can be described equivalently in terms of effective Lagrange (or Finsler) variables. This allows us to prove the second main result formulated in Theorem 3.1: Having chosen a generating Lagrange (Finsler) function $\mathcal{L}(x, y)$ (or $\mathcal{F}(x, y)$) on a (pseudo) Riemannian manifold, we can model this space as an almost Kähler geometry. Conventionally, the third main result is split into three (Theorems 5.1, 5.2 and 5.3) Fedosov's theorems for the normal distinguished connection in general relativity and its deformation quantization. We have introduced the normal Fedosov's distinguished operators, constructed the star product and computed the zero-degree cohomology coefficient for the almost Kähler models of (pseudo) Riemannian and Einstein spaces. The fourth result is given by Theorem 5.5, which states that a corresponding zero-degree cohomology coefficient also encodes the information about solutions of Einstein equations. This allows us to introduce, in Lagrange-

Finsler variables and using a related Chern–Cartan form, the quantum field equations corresponding to the classical gravitational equations in general relativity.

In this paper, we only concluded the first steps [1, 2, 3, 32] towards a consistent deformation quantization of gravity using the nonlinear connection formalism and the methods of Lagrange–Finsler geometry in Einstein gravity and generalizations. Many details of more complete constructions are still lacking. For example, we provided only the transformations suitable to implement deformation quantization methods but we have not discussed a de–quantization procedure and relevance to the semiclassical limit of such gravitational models. There are many unsolved problems pertaining canonical and quantum loop quantizations. It is very likely that non–holonomic geometry quantization methods of gravity and its relation to loop gravity, canonical and perturbative approaches, noncommutative generalizations and applications to modern cosmology and gravity physics will play an important role in future investigations.

Acknowledgement: The work was performed during a visit at Fields Institute. Author is grateful to Douglas Singleton and Andrea Arias de Gill from California State University at Fresno, USA, for substantial help on this paper.

References

- [1] S. Vacaru, Deformation Quantization of Almost Kähler Models and Lagrange–Finsler Spaces, *J. Math. Phys.* **48** (2007) 123509
- [2] S. Vacaru, Generalized Lagrange Transforms: Finsler Geometry Methods and Deformation Quantization of Gravity, *An. St. Univ. Al. I. Cuza din Iasi (S.N.), Matematica*, vol **LIII**, 2007, Supliment, 327-342
- [3] S. Vacaru, Deformation Quantization of Nonholonomic Almost Kähler Models and Einstein Gravity, *Phys. Lett. A* **372** (2008) 2949-2955
- [4] S. Vacaru, Branes and Quantization of an A–Model Complexification for Einstein Gravity in Almost Kähler Variables. *Int. J. Geom. Methods. Mod. Phys. (IJGMMP)* **6** (2009) 873-909
- [5] B. V. Fedosov, Deformation Quantization and Asymptotic Operator Representation, *Funktsional Anal. i Prilozhen.* **25** (1990) 1984–1994

- [6] B. V. Fedosov, A Simple Geometric Construction of Deformation Quantization, *J. Diff. Geom.* **40** (1994) 213–238
- [7] M. Kontsevich, Deformation Quantization of Poisson Manifolds, *Lett. Math. Phys.* **66** (2003) 157–216
- [8] M. Kontsevich, Operads and Motives in Deformation Quantization, *Lett. Math. Phys.* **48** (1999) 35–72
- [9] C. Rovelli, *Quantum Gravity* (Cambridge University Press, 2004)
- [10] A. Ashtekar and J. Lewandowski, Background Independent Quantum Gravity: A Status Report. *Class. Quant. Grav.* **21** (2004) R53
- [11] T. Thiemann, *Modern Canonical Quantum General Relativity* (Cambridge University Press, Cambridge, 2006)
- [12] L. Smolin, Quantum Gravity Faces Reality, *Phys. Today* **59** (2006) 44–48
- [13] J. H. Schwarz (Ed.), *Strings: the First 15 Years of Superstring Theory: Reprints & Commentary* (World Scientific, Singapore e. a., 1985)
- [14] P. Deligne, P. Etingof, D. S. Freed et al (eds.), *Quantum Fields and Strings: A Course for Mathematicians*, Vols 1 and 2, Institute for Advanced Study (American Mathematical Society, 1994)
- [15] J. Polchinski, *String Theory*, Vols 1 & 2 (Cambridge Univ. Press, 1998)
- [16] C. Misner, K. Thorne and J. Wheeler, *Gravitation* (Freeman, 1973)
- [17] F. A. Berezin, General Concept of Quantization, *Commun. Math. Phys.* **40** (1975) 153–174
- [18] F. Bayen, M. Flato, C. Fronsdal, A. Lichnerowicz and D. Sternheimer, Deformation Theory and Quantization. I. Deformations of Symplectic Structures, *Ann. Phys.* **111** (1978) 61–110
- [19] F. Bayen, M. Flato, C. Fronsdal, A. Lichnerowicz and D. Sternheimer, Deformation Theory and Quantization. II. Physical Applications, *Ann. Phys.* **111** (1978) 111–151
- [20] H. Garcia-Compean, J. Plebanski and M. Przanowski, The Geometry of Deformation Quantization and Self-Dual Gravity, in "Jerusalem 1997, Recent Developments in Theoretical and Experimental General

Relativity, Gravitation, and Relativistic Field Theories”, Pt. A (Marcel Grossman, Jerusalem, 1997), pp. 442–444

- [21] F. Antonsen, Deformation Quantization of Gravity, arXiv: gr-qc/ 9712012
- [22] H. Quevedo and J. Tafoya, Towards the Deformation Quantization of Linearized Gravity, *Gen. Rel. Grav.* **37** (2005) 2083–2092
- [23] C. Castro, W–Geometry from Fedosov’s Deformation Quantization, *J. Geom. Phys.* **33** (2000) 173–190
- [24] C. Castro, Moyal Deformations of Gravity via $SU(\infty)$ Gauge Theories, Branes and Topological Chern–Simons Matrix Models, *Gen. Rel. Grav.* **36** (2004) 2605–2635
- [25] Y. Cho, K. Soh, J. Yoon and W. Park, Gravitation as Gauge Theory of Diffeomorphism Group, *Phys. Lett. B* **286** (1992) 252–255
- [26] C. Castro, String and Membranes from Einstein Gravity, Matrix Models and W_∞ Gauge Theories, *Invertis J. Sci. Technol.* **2** (2008) 84–92
- [27] R. Miron and M. Anastasiei, Vector Bundles and Lagrange Spaces with Applications to Relativity (Geometry Balkan Press, Bukharest, 1997); translation from Romanian of (Editura Academiei Romane, 1987)
- [28] R. Miron and M. Anastasiei, The Geometry of Lagrange Spaces: Theory and Applications, FTPH no. 59 (Kluwer Academic Publishers, Dordrecht, Boston, London, 1994)
- [29] M. Anastasiei and S. Vacaru, Fedosov quantization of Lagrange-Finsler and Hamilton-Cartan spaces and Einstein gravity lifts on (co) tangent bundles, *J. Math. Phys.* **50** (2009) 013510
- [30] S. Vacaru, P. Stavrinou, E. Gaburov and D. Gonța, Clifford and Riemann- Finsler Structures in Geometric Mechanics and Gravity, Selected Works, Differential Geometry – Dynamical Systems, Monograph 7 (Geometry Balkan Press, 2006); www.mathem.pub.ro/dgds/mono/va-t.pdf
- [31] S. Vacaru, Finsler and Lagrange geometries in Einstein and string gravity, *Int. J. Geom. Methods. Mod. Phys. (IJGMMP)* **5** (2008) 473–511
- [32] F. Etayo, R. Santamaría and S. Vacaru, Lagrange-Fedosov Nonholonomic Manifolds, *J. Math. Phys.* **46** (2005) 032901

- [33] A. V. Karabegov and M. Schlichenmaier, Almost Kähler Deformation Quantization, *Lett. Math. Phys.* **57** (2001) 135–148
- [34] M. Matsumoto, *Foundations of Finsler Geometry and Special Publishers*, Dordrecht, Finsler Spaces (Kaisisha: Shigaken, 1986)
- [35] V. A. Oproiu, A Riemannian Structure in Lagrange Geometry, *Rendiconti Seminario Facoltà Scienze Università Cagliari*, **55** (1985) 1–20
- [36] P. O. Kazinski, S. L. Lyakhovich and A. A. Sharapov, Lagrange Structure and Quantization, *JHEP* 0507 (2005) 076
- [37] S. L. Lyakhovich and A. A. Sharapov, BRST Theory without Hamiltonian and Lagrangian, *JHEP* 0503 (2005) 011